

**DAVIS-MONTHAN AIR FORCE BASE  
OUTPATIENT FORMULARY  
Alphabetical Listing by Name**

This document is current as of May 28, 2020.  
The availability of formulary items is subject to change.

Davis-Monthan AFB Outpatient Formulary

| Alphabetical Drug Listing                                   | Available Dosage Forms   |
|---|--|
| Acetaminophen   | Caplet, oral: 325 mg<br>Suppository, rectal: 120 mg (12s); 650 mg (12s)<br>Suspension, oral: 160 mg/5 mL (5 mL)<br>Tablet, chewable, oral: 80 mg   |
| Acetaminophen and Codeine                                   | Solution, oral [C-V]: Acetaminophen 120 mg and codeine phosphate 12 mg per 5 mL (473 mL) [BCF]<br>Tablet, oral [C-III]: Acetaminophen 300 mg and codeine phosphate 30 mg [BCF]   |
| Acetaminophen and Tramadol                                  | Tablet, oral:<br>Ultracet: Acetaminophen 325 mg and tramadol hydrochloride 37.5 mg   |
| AcetaZOLAMIDE   | Capsule, sustained release, oral:<br>Diamox Sequels: 500 mg<br>Tablet, oral: 250 mg  |
| Acetic Acid (Otic)  | Solution, Otic:<br>Generic: 2% (15 mL)   |
| Acetic Acid, Propylene Glycol Diacetate, and Hydrocortisone | Solution, otic [drops]: Acetic acid 2%, propylene glycol diacetate 3%, and hydrocortisone 1% (10 mL)   |
| Acyclovir (Systemic)  | Capsule, oral: 200 mg [BCF]<br>Tablet, oral: 400 mg [BCF], 800 mg [BCF]  |
| Adalimumab  | Kit, Subcutaneous [preservative free]:<br>Humira: 40 mg/0.8 mL [contains polysorbate 80] [BCF]<br>Humira Pen: 40 mg/0.8 mL [contains polysorbate 80] [BCF]   |
| Adapalene   | Gel, topical:<br>Differin: 0.1% (45 g) [ethanol free]  |
| Albuterol   | Aerosol Solution, Inhalation:<br>ProAir HFA: 108 (90 Base) mcg/actuation (8.5 g) [BCF]<br>Solution, for nebulization: 0.5% [100 mg/20 mL] (1s)<br>Solution, for nebulization [preservative free]: 0.083% [2.5 mg/3 mL] (60s) [BCF]                                       |
| Alendronate   | Tablet, oral: 10 mg [BCF], 35 mg [BCF], 70 mg [BCF]  |
| Alendronate and Cholecalciferol                             | Tablet, oral:<br>Fosamax Plus D: 70/2800: Alendronate 70 mg and cholecalciferol 2800 int. units  |
| Alfuzosin   | Tablet, extended release, oral, as hydrochloride:<br>Uroxatral: 10 mg [BCF]  |
| Aliskiren   | Tablet, oral:<br>Tekturna: 150 mg, 300 mg  |
| Aliskiren and Hydrochlorothiazide                           | Tablet:<br>Tekturna HCT:<br>150/12.5: Aliskiren 150 mg and hydrochlorothiazide 12.5 mg<br>150/25: Aliskiren 150 mg and hydrochlorothiazide 25 mg<br>300/12.5: Aliskiren 300 mg and hydrochlorothiazide 12.5 mg<br>300/25: Aliskiren 300 mg and hydrochlorothiazide 25 mg |
| Allopurinol   | Tablet, oral: 100 mg [BCF], 300 mg [BCF]   |
| ALPRAZolam  | Tablet, oral: 0.5 mg<br>Xanax: 0.25 mg [scored]  |
| Alprostadil   | Injection, powder for reconstitution:<br>Caverject Impulse: 10 mcg [contains benzyl alcohol (in diluent), lactose 45.4 mg; cartridge; supplied with diluent]<br>Caverject: 20 mcg, 40 mcg [contains benzyl alcohol (in diluent), lactose 172 mg; supplied with diluent]  |
| Aluminum Chloride Hexahydrate                               | Solution, topical:<br>Drysol: 20% (35 mL) [contains ethanol 93%]   |
| Aluminum Hydroxide, Magnesium Hydroxide, and Simethicone    | Liquid, oral:<br>Maalox Advanced Regular Strength: Aluminum hydroxide 200 mg, magnesium hydroxide 200 mg, and simethicone 20 mg per 5 mL (355 mL) [contains magnesium 75 mg/5 mL, potassium 5 mg/5 mL, propylene glycol; mint flavor]                                    |

Davis-Monthan AFB Outpatient Formulary

|                             |   |
|-----------------------------|---|
| Amantadine                  | Capsule, oral, as hydrochloride: 100 mg [BCF]<br>Syrup, oral, as hydrochloride: 50 mg/5 mL (473 mL) [BCF]   |
| Amiodarone                  | Tablet, oral, as hydrochloride:<br>Pacerone: 200 mg [scored] [BCF]  |
| Amitriptyline               | Tablet, oral, as hydrochloride: 10 mg [BCF], 25 mg [BCF], 50 mg [BCF], 75 mg [BCF], 100 mg  |
| AmLODIPine                  | Tablet, oral: 2.5 mg [BCF], 5 mg [BCF], 10 mg [BCF]   |
| Amlodipine and Benazepril   | Capsule, oral:<br>Lotrel: 2.5/10: Amlodipine 2.5 mg and benazepril hydrochloride 10 mg [BCF]<br>Lotrel: 5/10: Amlodipine 5 mg and benazepril hydrochloride 10 mg [BCF]<br>Lotrel: 5/20: Amlodipine 5 mg and benazepril hydrochloride 20 mg [BCF]<br>Lotrel: 5/40: Amlodipine 5 mg and benazepril hydrochloride 40 mg [BCF]<br>Lotrel: 10/20: Amlodipine 10 mg and benazepril hydrochloride 20 mg [BCF]<br>Lotrel: 10/40: Amlodipine 10 mg and benazepril hydrochloride 40 mg [BCF]  |
| Amoxicillin                 | Capsule, oral: 250 mg [BCF], 500 mg [BCF]<br>Powder for suspension, oral: 125 mg/5 mL (150 mL); 250 mg/5 mL (150 mL) [BCF]; 400 mg/5 mL (100 mL) [BCF]<br>Tablet, oral: 875 mg<br>Tablet, chewable, oral: 250 mg  |
| Amoxicillin and Clavulanate | Powder for suspension, oral:<br>200: Amoxicillin 200 mg and clavulanate potassium 28.5 mg per 5 mL (100 mL) [BCF]<br>400: Amoxicillin 400 mg and clavulanate potassium 57 mg per 5 mL (100 mL) [BCF]<br>600: Amoxicillin 600 mg and clavulanate potassium 42.9 mg per 5 mL (125 mL) [BCF]<br>Tablet, oral:<br>250: Amoxicillin 250 mg and clavulanate potassium 125 mg [BCF]<br>500: Amoxicillin 500 mg and clavulanate potassium 125 mg [BCF]<br>875: Amoxicillin 875 mg and clavulanate potassium 125 mg [BCF]<br>Tablet, chewable, oral:<br>200: Amoxicillin 200 mg and clavulanate potassium 28.5 mg [contains phenylalanine]<br>400: Amoxicillin 400 mg and clavulanate potassium 57 mg [contains phenylalanine] |
| Apixaban                    | Tablet, Oral:<br>Eliquis: 2.5 mg [BCF], 5 mg [BCF]  |
| ARIPiprazole                | Tablet, oral:<br>Abilify: 2 mg, 5 mg, 10 mg, 15 mg, 30 mg   |
| Artificial Tears            | Solution, Ophthalmic:<br>HypoTears: Polyvinyl alcohol 1% and polyethylene glycol 400 1% (30 mL) [contains benzalkonium chloride]<br>Systane: Polyethylene glycol 400 0.4% and propylene glycol 0.3% (5 mL, 15 mL, 30 mL)  |
| Aspirin                     | Tablet, enteric coated, oral: 325 mg<br>Ecotrin Low Strength: 81 mg   |
| Atenolol                    | Tablet, oral: 25 mg [BCF], 50 mg [BCF], 100 mg [BCF]  |
| AtoMOXetine                 | Capsule, oral:<br>Strattera: 10 mg, 18 mg, 25 mg, 40 mg, 60 mg, 80 mg, 100 mg   |
| AtorvaSTATin                | Tablet, oral:<br>Lipitor: 10 mg [BCF], 20 mg [BCF], 40 mg [BCF], 80 mg [BCF]  |
| Atovaquone and Proguanil    | Tablet, oral:<br>Generic: Atovaquone 250 mg and proguanil hydrochloride 100 mg<br>Tablet, oral [pediatric]:<br>Generic: Atovaquone 62.5 mg and proguanil hydrochloride 25 mg  |
| Atropine (Ophthalmic)       | Solution, ophthalmic, as sulfate [drops]: 1% (15 mL)  |
| AzaTHIOprine                | Tablet, oral:<br>Imuran: 50 mg [scored]   |
| Azelastine (Nasal)          | Solution, intranasal, as hydrochloride [spray]:<br>Astelin: 0.1% [137 mcg/spray] (30 mL) [contains benzalkonium chloride; 200 metered sprays]   |

Davis-Monthan AFB Outpatient Formulary

|  |  |
|--|--|
| Azithromycin (Ophthalmic)                          | Solution, ophthalmic [drops]:<br>AzaSite: 1% (2.5 mL) [contains benzalkonium chloride]   |
| Azithromycin (Systemic)                            | Powder for suspension, oral, as dihydrate [strength expressed as base]:<br>Zithromax: 100 mg/5 mL (15 mL) [contains sodium 3.7 mg/5 mL; cherry-cr me de vanilla-banana flavor] [BCF]<br>Zithromax: 200 mg/5 mL (15 mL, 30 mL) [contains sodium 7.4 mg/5 mL; cherry-cr me de vanilla-banana flavor]<br>Tablet, oral, as dihydrate [strength expressed as base]:<br>Zithromax: 250 mg [contains sodium 0.9 mg/tablet] [BCF]<br>Tablet, oral, as monohydrate [strength expressed as base]: 250 mg [BCF] |
| Bacitracin and Polymyxin B (Ophthalmic)            | Ointment, ophthalmic: Bacitracin 500 units and polymyxin B 10,000 units per g (3.5 g)  |
| Bacitracin, Neomycin, and Polymyxin B (Ophthalmic) | Ointment, ophthalmic: Bacitracin 400 units, neomycin 3.5 mg, and polymyxin B 10,000 units per g (3.5 g) [BCF]  |
| Bacitracin (Topical)                               | Ointment, topical, as zinc [strength expressed as base]: 500 units/g (30 g)  |
| Baclofen   | Tablet, oral: 10 mg  |
| Benazepril   | Tablet, oral, as hydrochloride: 5 mg, 10 mg, 20 mg, 40 mg  |
| Benazepril and Hydrochlorothiazide                 | Tablet, oral:<br>5/6.25: Benazepril hydrochloride 5 mg and hydrochlorothiazide 6.25 mg<br>10/12.5: Benazepril hydrochloride 10 mg and hydrochlorothiazide 12.5 mg<br>20/12.5: Benazepril hydrochloride 20 mg and hydrochlorothiazide 12.5 mg<br>20/25: Benazepril hydrochloride 20 mg and hydrochlorothiazide 25 mg  |
| Benzocaine   | Lozenge, oral:<br>Cepacol Sore Throat: 15 mg (16s) [contains cetylpyridinium, menthol]   |
| Benzonatate  | Capsule, softgel, oral: 100 mg, 200 mg   |
| Benzoyl Peroxide                                   | Gel, External:<br>Generic: 5% (42.5 g)<br>Liquid, External:<br>Benzoyl Peroxide Wash: 5% (148 g, 237 g)<br>PanOxyl Wash: 10% (156 g) [contains alcohol, usp, methylparaben]  |
| Benzotropine                                       | Tablet, oral, as mesylate: 0.5 mg [BCF], 1 mg [BCF], 2 mg [BCF]  |
| Betamethasone and Clotrimazole                     | Cream, topical: Betamethasone dipropionate 0.05% (base) and clotrimazole 1% (15 g)   |
| Betamethasone (Topical)                            | Cream, topical, as dipropionate [strength expressed as base]: 0.05% (15 g)<br>Lotion, topical, as dipropionate [strength expressed as base]: 0.05% (60 mL)<br>Ointment, topical, as dipropionate [strength expressed as base]: 0.05% (15 g)  |
| Betaxolol (Ophthalmic)                             | Suspension, ophthalmic [drops]:<br>Betoptic S: 0.25% (10 mL) [contains benzalkonium chloride]  |
| Bethanechol  | Tablet, oral, as chloride: 50 mg<br>Urecholine: 25 mg [scored]   |
| Bicalutamide                                       | Tablet, Oral:<br>Generic: 50 mg [BCF]  |
| Bisacodyl  | Suppository, rectal:<br>Biscolax: 10 mg (100s)<br>Tablet, enteric coated, oral: 5 mg   |
| Bismuth Subsalicylate                              | Tablet, chewable, oral, as subsalicylate:<br>Peptic Relief: 262 mg   |
| Brimonidine (Ophthalmic)                           | Solution, ophthalmic, as tartrate [drops]: 0.2% (10 mL) [BCF]<br>Alphagan P: 0.1% (10 mL); 0.15% (10 mL) [contains Purite] [BCF]   |
| Bromfenac  | Solution, ophthalmic [drops]:<br>Prolensa: 0.07% (3 mL) [contains benzalkonium chloride, edetate disodium, sodium sulfite]   |

Davis-Monthan AFB Outpatient Formulary

|   |   |
|---|---|
| Budesonide and Formoterol               | Aerosol, for oral inhalation:<br>Symbicort: 80/4.5: Budesonide 80 mcg and formoterol fumarate dehydrate 4.5 mcg per actuation (10.2 g) [120 metered inhalations]<br>Symbicort: 160/4.5: Budesonide 160 mcg and formoterol fumarate dehydrate 4.5 mcg per actuation (10.2 g) [120 metered inhalations]   |
| Budesonide (Oral Inhalation)            | Suspension, Inhalation:<br>Generic: 0.25 mg/2 mL (2 mL); 0.5 mg/2 mL (2 mL); 1 mg/2 mL (2 mL)   |
| BuPROPion                               | Tablet, oral, as hydrochloride:<br>Wellbutrin: 75 mg [BCF], 100 mg [BCF]<br>Tablet, extended release, oral, as hydrochloride: 150 mg [generic for Wellbutrin SR] [BCF]; 200 mg [generic for Wellbutrin SR]<br>Wellbutrin XL: 150 mg [BCF], 300 mg [BCF]<br>Tablet, sustained release, oral, as hydrochloride:<br>Wellbutrin SR: 100 mg [BCF]<br>Zyban: 150 mg [BCF] |
| BusPIRone                               | Tablet, oral, as hydrochloride: 10 mg [BCF], 15 mg [BCF]  |
| Butalbital, Acetaminophen, and Caffeine | Tablet, oral:<br>Fioricet: Butalbital 50 mg, acetaminophen 325 mg, and caffeine 40 mg [BCF]   |
| Calcipotriene                           | Cream, topical:<br>Dovonex: 0.005% (60 g)   |
| Calcitriol (Systemic)                   | Capsule, softgel, oral:<br>Rocaltrol: 0.25 mcg [contains coconut oil]   |
| Capsaicin                               | Cream, topical:<br>Trixaicin: 0.025% (60 g) [contains benzyl alcohol]<br>Trixaicin HP: 0.075% (60 g) [contains benzyl alcohol]  |
| Captopril                               | Tablet, oral: 25 mg [BCF], 50 mg [BCF], 100 mg [BCF]  |
| CarBAMazepine                           | Suspension, oral:<br>TEGretol: 100 mg/5 mL (450 mL) [contains propylene glycol; citrus-vanilla flavor] [BCF]<br>Tablet, oral: 200 mg [BCF]<br>Tablet, chewable, oral:<br>TEGretol: 100 mg [scored] [BCF]<br>Tablet, extended release, oral:<br>TEGretol-XR: 100 mg, 200 mg  |
| Carbidopa and Levodopa                  | Tablet, oral: 25/100: Carbidopa 25 mg and levodopa 100 mg [BCF]<br>Sinemet: 10/100: Carbidopa 10 mg and levodopa 100 mg [scored] [BCF]<br>Sinemet: 25/250: Carbidopa 25 mg and levodopa 250 mg [scored] [BCF]<br>Tablet, extended release, oral:<br>25/100: Carbidopa 25 mg and levodopa 100 mg<br>50/200: Carbidopa 50 mg and levodopa 200 mg                      |
| Carboxymethylcellulose                  | Solution, ophthalmic, as sodium [drops, preservative free]:<br>Refresh Plus: 0.5% (0.4 mL)  |
| Carteolol (Ophthalmic)                  | Solution, ophthalmic, as hydrochloride [drops]: 1% (10 mL) [contains benzalkonium chloride]   |
| Carvedilol                              | Tablet, oral: 3.125 mg [BCF], 6.25 mg [BCF], 12.5 mg [BCF], 25 mg [BCF]   |
| Cefdinir                                | Capsule, oral: 300 mg<br>Powder for suspension, oral: 125 mg/5 mL (60 mL, 100 mL)   |
| Cefprozil                               | Powder for suspension, oral, as anhydrous: 250 mg/5 mL (100 mL)<br>Tablet, oral, as anhydrous: 250 mg   |
| Celecoxib                               | Capsule, oral:<br>CeleBREX: 100 mg, 200 mg  |
| Cephalexin                              | Capsule, oral: 500 mg [BCF]<br>Keflex: 250 mg [BCF]<br>Powder for suspension, oral: 125 mg/5 mL (100 mL); 250 mg/5 mL (200 mL) [BCF]  |
| Cetirizine (Systemic)                   | Syrup, oral, as hydrochloride: 5 mg/5 mL (120 mL)<br>Tablet, oral, as hydrochloride: 5 mg, 10 mg  |

Davis-Monthan AFB Outpatient Formulary

|                                  |  |
|----------------------------------|--|
| Chlorhexidine Gluconate (Oral)   | Solution, Mouth/Throat:<br>Periogard: 0.12% (473 mL) [mint flavor] [BCF]<br>Generic: 0.12% (473 mL) [BCF]  |
| Chlorpheniramine                 | Tablet, oral, as maleate:<br>Aller-chlor: 4 mg [scored]  |
| Chlorthalidone                   | Tablet, oral: 25 mg [BCF], 50 mg [BCF]   |
| Cholecalciferol                  | Solution, Oral [drops]:<br>Generic: 400 units/mL (50 mL)<br>Tablet, Oral:<br>Generic: 1000 units   |
| Ciclopirox                       | Solution, topical [nail lacquer]:<br>Penlac: 8% (6.6 mL) [contains isopropyl alcohol]  |
| Cimetidine                       | Tablet, oral: 400 mg   |
| Ciprofloxacin and Dexamethasone  | Suspension, otic:<br>Ciprodex: Ciprofloxacin 0.3% and dexamethasone 0.1% (7.5 mL) [contains benzalkonium chloride]   |
| Ciprofloxacin (Ophthalmic)       | Solution, ophthalmic, as hydrochloride [drops]: 3.5 mg/mL (5 mL) [equivalent to ciprofloxacin base 0.3%]   |
| Ciprofloxacin (Systemic)         | Tablet, oral, as hydrochloride [strength expressed as base]: 250 mg [BCF], 500 mg [BCF], 750 mg  |
| Citalopram                       | Solution, oral: 10 mg/5 mL (240 mL) [BCF]<br>Tablet, oral: 10 mg [BCF]<br>CeleXA: 20 mg [scored] [BCF], 40 mg [scored] [BCF]   |
| Clarithromycin                   | Granules for suspension, oral:<br>Biaxin: 125 mg/5 mL (100 mL); 250 mg/5 mL (100 mL) [fruit-punch flavor]<br>Tablet, oral:<br>Biaxin: 250 mg, 500 mg   |
| Clindamycin and Benzoyl Peroxide | Gel, External:<br>Clindamycin phosphate 1% and benzoyl peroxide 5% (50 g)<br>Clindamycin phosphate 1.2% and benzoyl peroxide 5% (45 g) [BCF]   |
| Clindamycin (Systemic)           | Capsule, oral, as hydrochloride [strength expressed as base]:<br>Cleocin HCl: 150 mg [contains tartrazine] [BCF]<br>Granules for solution, oral, as palmitate hydrochloride [strength expressed as base]:<br>Cleocin Pediatric: 75 mg/5 mL (100 mL) [cherry flavor]  |
| Clindamycin (Topical)            | Cream, vaginal, as phosphate [strength expressed as base]:<br>Cleocin: 2% (40 g) [contains benzyl alcohol, mineral oil] [BCF]<br>Gel, topical, as phosphate [strength expressed as base]:<br>Cleocin T: 1% (30 g)<br>Lotion, topical, as phosphate [strength expressed as base]: 1% (60 mL)<br>Solution, topical, as phosphate [strength expressed as base]:<br>Cleocin T: 1% (60 mL) [contains isopropyl alcohol 50%] [BCF] |
| Clobetasol                       | Cream, topical, as propionate: 0.05% (30 g)<br>Gel, topical, as propionate: 0.05% (30 g)<br>Ointment, topical, as propionate:<br>Temovate: 0.05% (30 g)<br>Solution, topical, as propionate [for scalp application]: 0.05% (25 mL)   |
| ClomiPHENE                       | Tablet, oral, as citrate:<br>Clomid: 50 mg [scored]  |
| Clonazepam                       | Tablet, oral: 0.5 mg [BCF], 1 mg   |
| CloNIDine                        | Tablet, oral, as hydrochloride: 0.1 mg [BCF], 0.2 mg [BCF]   |
| Clopidogrel                      | Tablet, oral:<br>Plavix: 75 mg [BCF]   |

Davis-Monthan AFB Outpatient Formulary

|                                   |   |
|-----------------------------------|---|
| Clotrimazole (Topical)            | Cream, topical: 1% (15 g)<br>Cream, vaginal:<br>Gyne-Lotrimin 7: 1% (45 g) [contains benzyl alcohol]<br>Solution, topical: 1% (10 mL) [BCF]   |
| Colchicine                        | Tablet, oral:<br>Colcrys: 0.6 mg [scored]   |
| Colestipol                        | Tablet, oral, as hydrochloride: 1 g   |
| Cortisone                         | Tablet, oral, as acetate: 25 mg   |
| Cromolyn (Ophthalmic)             | Solution, ophthalmic, as sodium [drops]: 4% (10 mL)   |
| Cyanocobalamin                    | Injection, solution: 1000 mcg/mL (1 mL)   |
| Cyclobenzaprine                   | Tablet, oral, as hydrochloride: 5 mg, 10 mg [BCF]   |
| Cyclopentolate                    | Solution, Ophthalmic, as hydrochloride:<br>Generic: 1% (2 mL, 15 mL)  |
| CycloSPORINE (Ophthalmic)         | Emulsion, ophthalmic [drops, preservative free]:<br>Restasis: 0.05% (0.4 mL) [contains castor oil]  |
| CycloSPORINE (Systemic)           | Capsule, oral [modified]: 25 mg, 100 mg   |
| Cyproheptadine                    | Tablet, oral, as hydrochloride: 4 mg [BCF]  |
| Dabigatran Etexilate              | Capsule, oral:<br>Pradaxa: 75 mg, 150 mg  |
| Desonide                          | Cream, topical: 0.05% (15 g)<br>Ointment, topical: 0.05% (15 g)   |
| DexAMETHasone (Systemic)          | Tablet, oral: 0.5 mg, 4 mg  |
| Dextroamphetamine                 | Capsule, extended release, oral, as sulfate: 10 mg<br>Tablet, oral, as sulfate: 5 mg  |
| Dextroamphetamine and Amphetamine | Capsule, extended release, oral:<br>Adderall XR: 5 mg [dextroamphetamine sulfate 1.25 mg, dextroamphetamine saccharate 1.25 mg, amphetamine aspartate monohydrate 1.25 mg, amphetamine sulfate 1.25 mg (equivalent to amphetamine base 3.1 mg)] [BCF]<br>Adderall XR: 10 mg [dextroamphetamine sulfate 2.5 mg, dextroamphetamine saccharate 2.5 mg, amphetamine aspartate monohydrate 2.5 mg, amphetamine sulfate 2.5 mg (equivalent to amphetamine base 6.3 mg)] [BCF]<br>Adderall XR: 15 mg [dextroamphetamine sulfate 3.75 mg, dextroamphetamine saccharate 3.75 mg, amphetamine aspartate monohydrate 3.75 mg, amphetamine sulfate 3.75 mg (equivalent to amphetamine base 9.4 mg)] [BCF]<br>Adderall XR: 20 mg [dextroamphetamine sulfate 5 mg, dextroamphetamine saccharate 5 mg, amphetamine aspartate monohydrate 5 mg, amphetamine sulfate 5 mg (equivalent to amphetamine base 12.5 mg)] [BCF]<br>Adderall XR: 25 mg [dextroamphetamine sulfate 6.25 mg, dextroamphetamine saccharate 6.25 mg, amphetamine aspartate monohydrate 6.25 mg, amphetamine sulfate 6.25 mg (equivalent to amphetamine base 15.6 mg)] [BCF]<br>Adderall XR: 30 mg [dextroamphetamine sulfate 7.5 mg, dextroamphetamine saccharate 7.5 mg, amphetamine aspartate monohydrate 7.5 mg, amphetamine sulfate 7.5 mg (equivalent to amphetamine base 18.8 mg)] [BCF]<br>Tablet, oral:<br>Adderall: 5 mg [scored; dextroamphetamine sulfate 1.25 mg, dextroamphetamine saccharate 1.25 mg, amphetamine aspartate monohydrate 1.25 mg, amphetamine sulfate 1.25 mg (equivalent to amphetamine base 3.13 mg)]<br>Adderall: 10 mg [scored; dextroamphetamine sulfate 2.5 mg, dextroamphetamine saccharate 2.5 mg, amphetamine aspartate monohydrate 2.5 mg, amphetamine sulfate 2.5 mg (equivalent to amphetamine base 6.3 mg)]<br>Adderall: 20 mg [scored; dextroamphetamine sulfate 5 mg, dextroamphetamine saccharate 5 mg, amphetamine aspartate monohydrate 5 mg, amphetamine sulfate 5 mg (equivalent to amphetamine base 12.6 mg)]<br>Adderall: 30 mg [scored; dextroamphetamine sulfate 7.5 mg, dextroamphetamine saccharate 7.5 mg, amphetamine aspartate monohydrate 7.5 mg, amphetamine sulfate 7.5 mg (equivalent |

Davis-Monthan AFB Outpatient Formulary

|                            |  |
|----------------------------|--|
| Dextrose                   | Gel, oral:<br>Glucose 15: 40% (37.5 g) [dye free; lemon flavor; provides dextrose 15 g/tube]   |
| Diabetic Supplies          | Glucose Meter:<br>FreeStyle Lite Glucose Monitor<br>Precision Xtra Device<br>TRUE2go Blood Glucose Monitoring Kit<br>TRUEresult Blood Glucose Monitoring System<br>Lancet:<br>FreeStyle Lancets: 100s<br>Syringe and needle:<br>BD Syringe: (3 mL) 23 gauge x 1"<br>BD Syringe Ultra-Fine: (0.3 mL) 31 gauge x 5/16"<br>BD Syringe Ultra-Fine: (1 mL) 30 gauge x 1/2"<br>BD Syringe Ultra-Fine II: (0.5 mL) 31 gauge x 5/16"<br>BD Syringe Ultra-Fine II: (1 mL) 31 gauge x 5/16"<br>BD TB Syringe: (1 mL) 25 gauge x 5/8"<br>NovoFine Pen Needle (30 gauge - 8 mm)<br>Test Strips/Kit:<br>FreeStyle Test Strips: 100s<br>Precision Xtra Blood Glucose Test Strips [BCF]<br>Precision Xtra Blood Glucose In Vitro Strip (100s)<br>TRUEtest In Vitro Strip (100s) |
| DiazePAM                   | Tablet, oral: 2 mg, 5 mg [BCF]   |
| Diclofenac (Systemic)      | Tablet, delayed release, enteric coated, oral, as sodium: 50 mg, 75 mg   |
| Diclofenac (Topical)       | Gel, topical, as sodium:<br>Voltaren Gel: 1% (100 g) [contains isopropyl alcohol]  |
| Dicloxacillin              | Capsule, oral: 250 mg [BCF]  |
| Dicyclomine                | Capsule, oral, as hydrochloride: 10 mg [BCF]<br>Syrup, oral, as hydrochloride: 10 mg/5 mL (473 mL) [BCF]<br>Tablet, oral, as hydrochloride: 20 mg [BCF]  |
| Digoxin                    | Solution, oral: 50 mcg/mL (60 mL)<br>Tablet, oral:<br>Lanoxin: 125 mcg [BCF], 250 mcg [BCF] [scored]   |
| DiI TIAZem                 | Capsule, extended release, oral, as hydrochloride [once-daily dosing]: 360 mg [BCF]<br>Tiazac: 120 mg [BCF], 180 mg [BCF], 240 mg [BCF], 300 mg [BCF]<br>Tablet, oral, as hydrochloride:<br>Cardizem: 60 mg [scored]   |
| DiphenhydrAMINE (Systemic) | Capsule, oral, as hydrochloride: 25 mg, 50 mg<br>Solution, oral, as hydrochloride: 12.5 mg/5 mL (5 mL)<br>Q-dryl: 12.5 mg/5 mL (120 mL) [ethanol free; contains sodium 5 mg/5 mL, sodium benzoate; cherry flavor]  |
| Diphenoxylate and Atropine | Tablet, oral:<br>Lomotil: 2.5/0.025: Diphenoxylate hydrochloride 2.5 mg and atropine sulfate 0.025 mg  |
| Docusate                   | Capsule, softgel, oral, as sodium: 100 mg<br>Liquid, oral, as sodium:<br>Colace: 10 mg/mL (30 mL) [contains propylene glycol, sodium 1 mg/mL]  |
| Donepezil                  | Tablet, oral, as hydrochloride:<br>Aricept: 5 mg, 10 mg  |
| Dorzolamide                | Solution, ophthalmic [drops]:<br>Trusopt: 2% (10 mL) [contains benzalkonium chloride]  |
| Dorzolamide and Timolol    | Solution, ophthalmic [drops]: Dorzolamide hydrochloride 2% [strength expressed as base] and timolol maleate 0.5% [strength expressed as base] (10 mL)  |
| Doxazosin                  | Tablet, oral: 1 mg, 4 mg<br>Cardura: 2 mg, 8 mg [scored]   |

Davis-Monthan AFB Outpatient Formulary

|   |   |
|---|---|
| Doxepin (Systemic)                              | Capsule, oral: 10 mg [BCF], 25 mg [BCF], 50 mg [BCF], 75 mg [BCF]<br>Solution, oral [concentrate]: 10 mg/mL (120 mL) [BCF]  |
| Doxycycline                                     | Capsule, oral, as hyclate [strength expressed as base]: 50 mg, 100 mg [BCF]<br>Tablet, oral, as hyclate [strength expressed as base]: 20 mg, 100 mg [BCF]   |
| Dronedarone                                     | Tablet, oral:<br>Multaq: 400 mg   |
| DULoxetine                                      | Capsule Delayed Release Particles, Oral:<br>Generic: 20 mg, 30 mg, 60 mg  |
| Eletriptan                                      | Tablet, oral:<br>Relpax: 20 mg, 40 mg   |
| Emollients                                      | Cream, topical:<br>Hydrocerin: (454 g)<br>Vanicream: (453 g)  |
| Emtricitabine and Tenofovir Disoproxil Fumarate | Tablet, Oral:<br>Truvada: Emtricitabine 100 mg and tenofovir disoproxil fumarate 150 mg, Emtricitabine 133 mg and tenofovir disoproxil fumarate 200 mg, Emtricitabine 167 mg and tenofovir disoproxil fumarate 250 mg, Emtricitabine 200 mg and tenofovir disoproxil fumarate 300 mg [contains fd&c blue #2 aluminum lake]  |
| Enoxaparin                                      | Injection, solution, as sodium [preservative free]:<br>Lovenox: 30 mg/0.3 mL (0.3 mL) [BCF]; 40 mg/0.4 mL (0.4 mL) [BCF]; 60 mg/0.6 mL (0.6 mL) [BCF]; 80 mg/0.8 mL (0.8 mL) [BCF]; 100 mg/mL (1 mL) [BCF]; 120 mg/0.8 mL (0.8 mL) [BCF]; 150 mg/mL (1 mL) [BCF] [prefilled syringe]  |
| EPINEPHrine (Systemic)                          | Injection, solution:<br>EpiPen 2-Pak: 0.3 mg/0.3 mL (2 mL) [contains sodium metabisulfite; 1:1000 solution; delivers 0.3 mg per injection]<br>EpiPen Jr 2-Pak: 0.15 mg/0.3 mL (2 mL) [contains sodium metabisulfite; 1:2000 solution; delivers 0.15 mg per injection]<br>Twinject: 0.15 mg/0.15 mL (1.1 mL) [contains chlorobutanol, sodium bisulfite; 1:1000 solution; delivers 0.15 mg per injection]<br>Twinject: 0.3 mg/0.3 mL (1.1 mL) [contains chlorobutanol, sodium bisulfite; 1:1000 solution; delivers 0.3 mg per injection]  |
| Ergocalciferol                                  | Capsule, oral:<br>Drisdol: 50,000 units [contains soybean oil, tartrazine; 1.25 mg]   |
| Erythromycin (Ophthalmic)                       | Ointment, Ophthalmic:<br>Generic: 5 mg/g (3.5 g) [BCF]  |
| Erythromycin (Systemic)                         | Tablet, delayed release, enteric coated, oral, as base:<br>Ery-Tab: 250 mg [contains sodium 8.3 mg (0.4 mEq)/tablet] [BCF]  |
| Erythromycin (Topical)                          | Gel, topical: 2% (30 g) [BCF]<br>Solution, topical: 2% (60 mL) [BCF]  |
| Escitalopram                                    | Tablet, oral: 5 mg, 10 mg, 20 mg  |
| Estradiol (Systemic)                            | Patch Weekly, Transdermal:<br>Climara: 0.025 mg/24 hours (4s) [6.5 cm <sup>2</sup> , total estradiol 2.04 mg] [BCF]<br>Climara: 0.0375 mg/24 hours (4s) [9.375 cm <sup>2</sup> , total estradiol 2.85 mg] [BCF]<br>Climara: 0.05 mg/24 hours (4s) [12.5 cm <sup>2</sup> , total estradiol 3.8 mg] [BCF]<br>Climara: 0.075 mg/24 hours (4s) [18.75 cm <sup>2</sup> , total estradiol 5.7 mg] [BCF]<br>Climara: 0.1 mg/24 hours (4s) [25 cm <sup>2</sup> , total estradiol 7.6 mg] [BCF]<br>Patch Twice Weekly, Transdermal:<br>Vivelle-Dot: 0.025 mg/24 hr (8 ea)<br>Vivelle-Dot: 0.0375 mg/24 hours (24s) [3.75 cm <sup>2</sup> , total estradiol 0.585 mg]<br>Vivelle-Dot: 0.05 mg/24 hours (24s) [5 cm <sup>2</sup> , total estradiol 0.78 mg]<br>Vivelle-Dot 0.1 mg/24 hours (24s) [10 cm <sup>2</sup> , total estradiol 1.56 mg]<br>Tablet, Oral [micronized]:<br>Generic: 0.5 mg, 1 mg, 2 mg |
| Estradiol (Topical)                             | Cream, vaginal:<br>Estrace: 0.1 mg/g (42.5 g)   |

Davis-Monthan AFB Outpatient Formulary

|   |  |
|---|--|
| Estrogens (Conjugated/Equine) and Medroxyprogesterone | Tablet, oral:<br>Prempro: 0.3/1.5: Conjugated estrogens 0.3 mg and medroxyprogesterone acetate 1.5 mg [BCF]<br>Prempro: 0.45/1.5: Conjugated estrogens 0.45 mg and medroxyprogesterone acetate 1.5 mg [BCF]<br>Prempro: 0.625/2.5: Conjugated estrogens 0.625 mg and medroxyprogesterone acetate 2.5 mg [BCF]  |
| Estrogens (Conjugated/Equine, Systemic)               | Tablet, oral:<br>Premarin: 0.3 mg [BCF], 0.45 mg [BCF], 0.625 mg [BCF], 0.9 mg, 1.25 mg [BCF]  |
| Estrogens (Conjugated/Equine, Topical)                | Cream, vaginal:<br>Premarin: 0.625 mg/g (30 g) [BCF]   |
| Estrogens (Esterified) and Methyltestosterone         | Tablet, oral: Esterified estrogen 0.625 mg and methyltestosterone 1.25 mg  |
| Eszopiclone   | Tablet, oral:<br>Lunesta: 1 mg, 2 mg, 3 mg   |
| Ethambutol  | Tablet, oral, as hydrochloride:<br>Myambutol: 400 mg [scored] [BCF]  |
| Ethinyl Estradiol and Desogestrel                     | Tablet, oral [monophasic formulation]:<br>Ortho-Cept: Ethinyl estradiol 0.03 mg and desogestrel 0.15 mg [21 light orange tablets and 7 green inactive tablets] (28s)<br>Tablet, oral [triphasic formulation]:<br>Cyclessa:<br>Day 1-7: Ethinyl estradiol 0.025 mg and desogestrel 0.1 mg [7 light yellow tablets]<br>Day 8-14: Ethinyl estradiol 0.025 mg and desogestrel 0.125 mg [7 orange tablets]<br>Day 15-21: Ethinyl estradiol 0.025 mg and desogestrel 0.15 mg [7 red tablets]<br>Day 22-28: 7 green inactive tablets (28s)  |
| Ethinyl Estradiol and Drospirenone                    | Tablet, oral:<br>Yasmin: Ethinyl estradiol 0.03 mg and drospirenone 3 mg [21 yellow active tablets and 7 white inactive tablets] (28s) [BCF]<br>Yaz: Ethinyl estradiol 0.02 mg and drospirenone 3 mg [24 light pink active tablets and 4 white inactive tablets] (28s) [BCF]   |
| Ethinyl Estradiol and Etonogestrel                    | Ring, vaginal:<br>NuvaRing: Ethinyl estradiol 0.015 mg/day and etonogestrel 0.12 mg/day (3s) [3-week duration]   |
| Ethinyl Estradiol and Levonorgestrel                  | Tablet, oral [extended cycle regimen]:<br>Jolessa: Ethinyl estradiol 0.03 mg and levonorgestrel 0.15 mg (91s) [84 pink tablets and 7 white inactive tablets] [BCF]<br>Tablet, oral [low dose formulation]:<br>Sronyx: Ethinyl estradiol 0.02 mg and levonorgestrel 0.1 mg (28s) [21 white tablets and 7 peach inactive tablets] [BCF]<br>Tablet, oral [monophasic formulation]:<br>Nordette 28: Ethinyl estradiol 0.03 mg and levonorgestrel 0.15 mg (28s) [21 light orange tablets and 7 pink inactive tablets] [BCF]   |
| Ethinyl Estradiol and Norelgestromin                  | Patch, transdermal:<br>Ortho Evra: Ethinyl estradiol 0.75 mg and norelgestromin 6 mg [releases ethinyl estradiol 20 mcg and norelgestromin 150 mcg per day] (3s)   |
| Ethinyl Estradiol and Norethindrone                   | Tablet, oral:<br>Jinteli: Ethinyl estradiol 0.005 mg and norethindrone acetate 1 mg [white tablets] (90s)<br>Tablet, oral [monophasic formulation]:<br>Loestrin Fe 1/20: Ethinyl estradiol 0.02 mg and norethindrone acetate 1 mg [21 light yellow tablets] and ferrous fumarate 75 mg [7 brown tablets] (28s)<br>Loestrin Fe 1.5/30: Ethinyl estradiol 0.03 mg and norethindrone acetate 1.5 mg [21 pink tablets] and ferrous fumarate 75 mg [7 brown tablets] (28s)<br>Ortho-Novum 1/35: Ethinyl estradiol 0.035 mg and norethindrone 1 mg (28s) [21 peach tablets and 7 green inactive tablets] [BCF] |

Davis-Monthan AFB Outpatient Formulary

|                                    |  |
|------------------------------------|--|
| Ethinyl Estradiol and Norgestimate | Tablet, oral [monophasic formulation]:<br>Ortho-CyclenÂ®: Ethinyl estradiol 0.035 mg and norgestimate 0.25 mg [21 blue tablets and 7 dark green inactive tablets] (28s)<br>Tablet, oral [triphasic formulation]:<br>Ortho Tri-Cyclen:<br>Day 1-7: Ethinyl estradiol 0.035 mg and norgestimate 0.18 mg [7 white tablets]<br>Day 8-14: Ethinyl estradiol 0.035 mg and norgestimate 0.215 mg [7 light blue tablets]<br>Day 15-21: Ethinyl estradiol 0.035 mg and norgestimate 0.25 mg [7 blue tablets]<br>Day 22-28: 7 dark green inactive tablets (28s)<br>Ortho Tri-Cyclen Lo [BCF]:<br>Day 1-7: Ethinyl estradiol 0.025 mg and norgestimate 0.18 mg [7 white tablets]<br>Day 8-14: Ethinyl estradiol 0.025 mg and norgestimate 0.215 mg [7 light blue tablets]<br>Day 15-21: Ethinyl estradiol 0.025 mg and norgestimate 0.25 mg [7 dark blue tablets]<br>Day 22-28: 7 dark green inactive tablets |
| Ethinyl Estradiol and Norgestrel   | Tablet, oral [monophasic formulation]:<br>Low-Ogestrel: Ethinyl estradiol 0.03 mg and norgestrel 0.3 mg [21 white tablets and 7 peach inactive tablets] (28s)  |
| Exenatide                          | Auto-injector, Subcutaneous:<br>Bydureon BCise: 2 mg/0.85 mL (0.85 mL) [BCF]   |
| Ezetimibe                          | Tablet, oral:<br>Zetia: 10 mg  |
| Famotidine                         | Tablet, Oral:<br>Generic: 20 mg, 40 mg   |
| Felodipine                         | Tablet, extended release, oral: 2.5 mg, 5 mg, 10 mg  |
| Fenofibrate and Derivatives        | Capsule, oral [micronized]: 67 mg, 134 mg<br>Lofibra: 200 mg<br>Tablet, oral:<br>TriCor: 48 mg [BCF], 145 mg [BCF] [contains soybean lecithin]   |
| Ferrous Sulfate                    | Liquid, oral [drops]: 75 mg/mL (50 mL) [elemental iron 15 mg/mL]<br>Tablet, enteric coated, oral: 324 mg [elemental iron 65 mg]  |
| Fexofenadine                       | Tablet, oral, as hydrochloride: 60 mg, 180 mg<br>Allegra Children's Allergy: 30 mg   |
| Fexofenadine and Pseudoephedrine   | Tablet, extended release, oral:<br>Allegra-D 24 Hour: Fexofenadine hydrochloride 180 mg [immediate release] and pseudoephedrine hydrochloride 240 mg [extended release]  |
| Finasteride                        | Tablet, oral:<br>Proscar: 5 mg [BCF]   |
| Flecainide                         | Tablet, Oral, as acetate:<br>Generic: 50 mg, 100 mg, 150 mg  |
| Fluconazole                        | Powder for suspension, oral: 40 mg/mL (35 mL)<br>Tablet, oral: 50 mg, 100 mg, 150 mg [BCF], 200 mg   |
| Fludrocortisone                    | Tablet, oral, as acetate: 0.1 mg   |
| Fluocinolone (Topical)             | Oil, topical, as acetonide [scalp oil]:<br>Derma-Smoothe/FS: 0.01% (120 mL) [contains isopropyl alcohol, peanut oil]<br>Solution, topical, as acetonide: 0.01% (60 mL)   |
| Fluocinonide                       | Cream, anhydrous, emollient, topical: 0.05% (15 g) [BCF]<br>Cream, aqueous, emollient, topical: 0.05% (15 g)<br>Gel, topical: 0.05% (15 g, 60 g)<br>Ointment, topical: 0.05% (15 g)  |
| Fluoride                           | Cream, oral, as sodium [toothpaste]:<br>PreviDent 5000 Plus: 1.1% (51 g) [contains sodium benzoate; fruitastic flavor; equivalent to fluoride 2.5 mg/dose]   |
| Fluorometholone                    | Suspension, ophthalmic, as base [drops]:<br>FML: 0.1% (10 mL) [contains benzalkonium chloride]   |

Davis-Monthan AFB Outpatient Formulary

|                                    |  |
|------------------------------------|--|
| Fluorouracil (Topical)             | Cream, topical:<br>Carac: 0.5% (30 g)<br>Efudex: 5% (40 g)   |
| FLUoxetine                         | Capsule, oral: 20 mg [BCF], 40 mg [BCF]<br>PROzac: 10 mg [BCF]<br>Solution, oral: 20 mg/5 mL (120 mL) [BCF]  |
| Flurbiprofen (Ophthalmic)          | Solution, ophthalmic, as sodium [drops]:<br>Ocufen: 0.03% (2.5 mL)   |
| Fluticasone and Salmeterol         | Aerosol, for oral inhalation:<br>Advair HFA: 45/21: Fluticasone propionate 45 mcg and salmeterol 21 mcg per inhalation (12 g) [chlorofluorocarbon free; 120 metered actuations] [BCF]<br>Advair HFA: 115/21: Fluticasone propionate 115 mcg and salmeterol 21 mcg per inhalation (12 g) [chlorofluorocarbon free; 120 metered actuations] [BCF]<br>Advair HFA: 230/21: Fluticasone propionate 230 mcg and salmeterol 21 mcg per inhalation (12 g) [chlorofluorocarbon free; 120 metered actuations] [BCF]<br>Powder, for oral inhalation:<br>Advair Diskus: 100/50: Fluticasone propionate 100 mcg and salmeterol 50 mcg (14s, 60s) [contains lactose] [BCF]<br>Advair Diskus: 250/50: Fluticasone propionate 250 mcg and salmeterol 50 mcg (14s, 60s) [contains lactose] [BCF]<br>Advair Diskus: 500/50: Fluticasone propionate 500 mcg and salmeterol 50 mcg (14s, 60s) [contains lactose] [BCF] |
| Fluticasone (Nasal)                | Suspension, intranasal, as propionate [spray]:<br>Flonase: 50 mcg/inhalation (16 g) [contains benzalkonium chloride; 120 metered actuations] [BCF]   |
| Fluticasone (Oral Inhalation)      | Aerosol, for oral inhalation, as propionate:<br>Flovent HFA: 44 mcg/inhalation (10.6 g) [chlorofluorocarbon free; 120 metered actuations] [BCF]<br>Flovent HFA: 110 mcg/inhalation (12 g) [chlorofluorocarbon free; 120 metered actuations] [BCF]<br>Flovent HFA: 220 mcg/inhalation (12 g) [chlorofluorocarbon free; 120 metered actuations] [BCF]<br>Powder, for oral inhalation, as propionate:<br>Flovent Diskus: 50 mcg (60s) [contains lactose] [BCF]<br>Flovent Diskus: 100 mcg (60s) [contains lactose] [BCF]<br>Flovent Diskus: 250 mcg (60s) [contains lactose] [BCF]  |
| Fluticasone (Topical)              | Cream, topical, as propionate: 0.05% (30 g)  |
| Fluvoxamine                        | Tablet, oral, as maleate: 50 mg, 100 mg  |
| Folic Acid                         | Tablet, oral: 1 mg [BCF]   |
| Fosfomycin                         | Packet, Oral:<br>Monurol: 3 g (1 ea) [orange flavor]   |
| Fosinopril                         | Tablet, oral, as sodium: 10 mg, 20 mg, 40 mg   |
| Fosinopril and Hydrochlorothiazide | Tablet, oral: 10/12.5: Fosinopril sodium 10 mg and hydrochlorothiazide 12.5 mg; 20/12.5: Fosinopril sodium 20 mg and hydrochlorothiazide 12.5 mg   |
| Furosemide                         | Solution, oral: 10 mg/mL (120 mL) [BCF]<br>Tablet, oral: 40 mg [BCF]<br>Lasix: 20 mg [BCF]   |
| Gabapentin                         | Capsule, oral: 300 mg [BCF]<br>Neurontin: 100 mg [BCF] 400 mg [BCF]<br>Tablet, oral: 800 mg [BCF]<br>Neurontin: 600 mg [BCF]   |
| Ganciclovir (Ophthalmic)           | Gel, ophthalmic [drops]:<br>Zirgan: 0.15% (5 g) [contains benzalkonium chloride]   |
| Gemfibrozil                        | Tablet, oral: 600 mg [BCF]   |

Davis-Monthan AFB Outpatient Formulary

|                                     |  |
|-------------------------------------|--|
| Gentamicin (Ophthalmic)             | Ointment, ophthalmic:<br>Gentak: 0.3% (3.5 g) [BCF]<br>Solution, ophthalmic [drops]: 0.3% (5 mL, 15 mL) [BCF]  |
| Gentamicin (Topical)                | Cream, topical: 0.1% (15 g, 30 g)  |
| Glimepiride                         | Tablet, oral: 1 mg, 2 mg, 4 mg   |
| GlipiZIDE                           | Tablet, oral: 5 mg [BCF], 10 mg [BCF]<br>Tablet, extended release, oral: 2.5 mg, 5 mg, 10 mg   |
| Glucagon                            | Injection, powder for reconstitution:<br>GlucaGen Diagnostic Kit: 1 mg [contains lactose 107 mg; equivalent to 1 unit]   |
| GlyBURIDE                           | Tablet, oral: 1.25 mg [BCF], 2.5 mg [BCF], 5 mg [BCF]<br>Tablet, oral [micronized]: 6 mg [BCF]<br>Glynase PresTab: 3 mg [BCF]  |
| Glyburide and Metformin             | Tablet, oral:<br>Glucovance: 5/500: Glyburide 5 mg and metformin hydrochloride 500 mg, 2.5/500: Glyburide 2.5 mg and metformin hydrochloride 500 mg  |
| Glycerin                            | Suppository, rectal [pediatric]:<br>Sani-Supp: 82.5% (10s)   |
| Goserelin                           | Implant, subcutaneous:<br>Zoladex: 3.6 mg (1s) [1 month implant] [BCF]<br>Zoladex: 10.8 mg (1s) [3 month implant] [BCF]  |
| Griseofulvin                        | Suspension, oral [microsize]: 125 mg/5 mL (120 mL) [BCF]<br>Tablet, oral [ultramicrosize]:<br>Gris-PEG: 250 mg [BCF] [scored]  |
| GuaiFENesin                         | Syrup, oral: 100 mg/5 mL (118 mL)<br>Tablet, extended release, oral:<br>Mucinex: 600 mg  |
| Guaifenesin and Codeine             | Solution, oral: Guaifenesin 100 mg and codeine phosphate 10 mg per 5 mL (118 mL)   |
| Guaifenesin and Dextromethorphan    | Syrup, oral: Guaifenesin 100 mg and dextromethorphan hydrobromide 10 mg per 5 mL (118 mL)  |
| Guaifenesin and Pseudoephedrine     | Tablet, extended release, oral:<br>Mucinex D: Guaifenesin 600 mg and pseudoephedrine hydrochloride 60 mg   |
| GuanFACINE                          | Tablet, Oral:<br>Generic: 1 mg, 3 mg<br>Tablet Extended Release 24 Hour, Oral:<br>Generic: 1 mg, 2 mg  |
| Haloperidol                         | Solution, oral, as lactate [strength expressed as base, concentrate]: 2 mg/mL (120 mL)<br>Tablet, oral: 1 mg, 2 mg, 10 mg  |
| Homatropine                         | Solution, ophthalmic, as hydrobromide [drops]:<br>Homatropaire: 5% (5 mL) [contains benzalkonium chloride]<br>Isopto Homatropine: 2% (5 mL) [contains benzalkonium chloride]   |
| HydrALAZINE                         | Tablet, oral, as hydrochloride: 10 mg [BCF], 25 mg [BCF], 50 mg [BCF]  |
| HydroCHLORothiazide                 | Capsule, oral: 12.5 mg [BCF]<br>Tablet, oral: 25 mg [BCF], 50 mg [BCF]   |
| Hydrochlorothiazide and Triamterene | Tablet, oral: Hydrochlorothiazide 25 mg and triamterene 37.5 mg [BCF]; hydrochlorothiazide 50 mg and triamterene 75 mg [BCF]   |
| Hydrocodone and Acetaminophen       | Tablet, oral:<br>Hydrocodone bitartrate 5 mg and acetaminophen 500 mg<br>Hydrocodone bitartrate 5 mg and acetaminophen 325 mg<br>Hydrocodone bitartrate 7.5 mg and acetaminophen 325 mg<br>Hydrocodone bitartrate 7.5 mg and acetaminophen 750 mg<br>Hydrocodone bitartrate 10 mg and acetaminophen 325 mg |
| Hydrocortisone (Systemic)           | Tablet, oral, as base:<br>Generic: 5 mg  |

Davis-Monthan AFB Outpatient Formulary

|   |  |
|---|--|
| Hydrocortisone (Topical)                              | Cream, topical, as base: 1% (28.4 g)<br>Anusol-HC: 2.5% (30 g) [contains benzyl alcohol] [BCF]<br>Cream, topical, as valerate: 0.2% (15 g)<br>Ointment, topical, as base: 1% (30 g); 2.5% (20 g)<br>Ointment, topical, as valerate:<br>Westcort: 0.2% (60 g)<br>Suppository, rectal, as acetate: 25 mg (12s) [BCF]   |
| Hydroxychloroquine                                    | Tablet, oral, as sulfate: 200 mg [equivalent to 155 mg base]   |
| Hydroxyurea   | Capsule, oral:<br>Hydrea: 500 mg   |
| HydrOXYzine   | Capsule, oral, as pamoate:<br>Vistaril: 25 mg<br>Syrup, oral, as hydrochloride: 10 mg/5 mL (473 mL) [BCF]<br>Tablet, oral, as hydrochloride: 10 mg [BCF], 25 mg [BCF]  |
| Hyoscyamine   | Tablet, sublingual, as sulfate:<br>Levsin/SL: 0.125 mg   |
| Hyoscyamine, Atropine, Scopolamine, and Phenobarbital | Elixir, oral:<br>Donnatal: Hyoscyamine sulfate 0.1037 mg, atropine sulfate 0.0194 mg, scopolamine hydrobromide 0.0065 mg, and phenobarbital 16.2 mg per 5 mL (118 mL) [contains ethanol <23.8%; grape flavor]<br>Tablet, oral:<br>Donnatal: Hyoscyamine sulfate 0.1037 mg, atropine sulfate 0.0194 mg, scopolamine hydrobromide 0.0065 mg, and phenobarbital 16.2 mg |
| Ibandronate   | Tablet, oral: 150 mg   |
| Ibuprofen   | Suspension, oral: 100 mg/5 mL (120 mL)<br>Tablet, oral: 400 mg [BCF], 600 mg [BCF], 800 mg [BCF]   |
| Imipramine  | Capsule, oral, as pamoate:<br>Tofranil-PM: 75 mg [BCF]<br>Tablet, oral, as hydrochloride: 25 mg [BCF]<br>Tofranil: 10 mg [BCF], 50 mg [BCF]  |
| Imiquimod   | Cream, topical:<br>Aldara: 5% (12s) [contains benzyl alcohol; 0.25 g/packet]   |
| Indomethacin  | Capsule, oral: 25 mg [BCF], 50 mg [BCF]  |
| Inhalation Devices                                    | Holding chamber with mask:<br>AeroChamber Plus: Small, medium, large<br>Inhalation kit:<br>InspirEase  |
| Insulin Aspart  | Injection, solution:<br>NovoLOG: 100 units/mL (10 mL) [vial] [BCF]<br>NovoLOG FlexPen: 100 units/mL (3 mL) [BCF]<br>NovoLOG Penfill: 100 units/mL (3 mL) [cartridge] [BCF]   |
| Insulin Aspart Protamine and Insulin Aspart           | Injection, suspension:<br>NovoLOG Mix 70/30: Insulin aspart protamine suspension 70% [intermediate acting] and insulin aspart solution 30% [rapid acting]: 100 units/mL (10 mL) [BCF]<br>NovoLOG Mix 70/30 FlexPen: Insulin aspart protamine suspension 70% [intermediate acting] and insulin aspart solution 30% [rapid-acting] (3 mL) [BCF]                        |
| Insulin Detemir                                       | Injection, solution:<br>Levemir: 100 units/mL (10 mL)  |
| Insulin Glargine                                      | Injection, solution:<br>Lantus: 100 units/mL (10 mL) [vial] [BCF]<br>Lantus Solostar: 100 units/mL (3 mL) [prefilled pen] [BCF]  |
| Insulin NPH   | Injection, suspension:<br>NovoLIN N: 100 units/mL (10 mL) [vial] [BCF]   |

Davis-Monthan AFB Outpatient Formulary

|                                 |   |
|---------------------------------|---|
| Insulin NPH and Insulin Regular | Injection, suspension:<br>NovoLIN 70/30: Insulin NPH suspension 70% [intermediate acting] and insulin regular solution 30% [short acting]: 100 units/mL (10 mL) [vial] [BCF]  |
| Insulin Regular                 | Injection, solution:<br>NovoLIN R: 100 units/mL (10 mL) [BCF]   |
| Ipratropium and Albuterol       | Aerosol, for oral inhalation:<br>Combivent: Ipratropium bromide 18 mcg and albuterol (base) 90 mcg per inhalation (14.7 g) [contains chlorofluorocarbon, soya lecithin; 200 metered actuations] [BCF]<br>Solution, for nebulization: Ipratropium bromide 0.5 mg and albuterol (base) 2.5 mg per 3 mL (60s)<br>Solution, for oral inhalation [spray]:<br>Combivent Respimat: Ipratropium bromide 20 mcg and albuterol (base) 100 mcg per inhalation (4 g) [contains benzalkonium chloride; 120 metered actuations] |
| Ipratropium (Nasal)             | Solution, intranasal, as bromide [spray]: 0.03% (30 mL); 0.06% (15 mL)  |
| Ipratropium (Oral Inhalation)   | Aerosol, for oral inhalation, as bromide:<br>Atrovent HFA: 17 mcg/actuation (12.9 g) [chlorofluorocarbon free; 200 metered actuations] [BCF]<br>Solution, for nebulization, as bromide [preservative free]: 0.02% [500 mcg/2.5 mL] ( 30s)   |
| Isoniazid                       | Solution, oral: 50 mg/5 mL (473 mL)<br>Tablet, oral: 300 mg [BCF]   |
| Isosorbide Dinitrate            | Capsule, sustained release, oral:<br>Dilatrate-SR: 40 mg<br>Tablet, oral: 10 mg [BCF], 20 mg [BCF], 30 mg [BCF]   |
| Isosorbide Mononitrate          | Tablet, extended release, oral: 30 mg [BCF]<br>Imdur: 60 mg [scored] [BCF]  |
| ISOTretinoin (Systemic)         | Capsule, Oral:<br>Generic: 10 mg, 20 mg, 30 mg, 40 mg   |
| Ketoconazole (Systemic)         | Tablet, oral: 200 mg  |
| Ketoconazole (Topical)          | Cream, topical: 2% (30 g)<br>Shampoo, topical:<br>Nizoral: 2% (120 mL)  |
| Ketorolac (Ophthalmic)          | Solution, ophthalmic, as tromethamine [drops]: 0.4% (5 mL); 0.5% (5 mL)   |
| Ketotifen (Ophthalmic)          | Solution, ophthalmic [drops]:<br>Zaditor: 0.025% (5 mL) [contains benzalkonium chloride]  |
| Lactulose                       | Solution, oral: 10 g/15 mL (946 mL) [BCF]   |
| Lamotrigine                     | Tablet, oral: 25 mg, 100 mg, 200 mg   |
| Latanoprost                     | Solution, ophthalmic [drops]:<br>Xalatan: 0.005% (2.5 mL) [contains benzalkonium chloride] [BCF]  |
| Leuprolide                      | Injection, powder for reconstitution, as acetate [depot formulation, preservative free]:<br>Lupron Depot: 3.75 mg (monthly) [contains polylactide-co-glycolide, polysorbate 80]   |
| Levetiracetam                   | Tablet, oral: 250 mg, 500 mg, 750 mg  |
| Levofloxacin (Systemic)         | Tablet, oral:<br>Levaquin: 250 mg [BCF], 500 mg [BCF], 750 mg [BCF]   |
| Levonorgestrel (Systemic)       | Tablet, Oral:<br>Plan B: 0.75 mg [BCF]  |
| Levothyroxine                   | Tablet, oral, as sodium:<br>Synthroid: 25 mcg [BCF], 50 mcg [scored; dye free] [BCF], 75 mcg [BCF], 88 mcg [BCF], 100 mcg [BCF], 112 mcg [BCF], 125 mcg [BCF], 137 mcg [BCF], 150 mcg [BCF], 175 mcg [BCF], 200 mcg [BCF]   |
| Lidocaine (Topical)             | Patch, topical:<br>Lidoderm: 5% (30s)<br>Solution, viscous, oral topical, as hydrochloride: 2% [20 mg/mL] (100 mL)  |
| Liothyronine                    | Tablet, oral:<br>Cytomel: 5 mcg<br>Cytomel: 25 mcg [scored]   |

Davis-Monthan AFB Outpatient Formulary

|                                    |  |
|------------------------------------|--|
| Lisinopril                         | Tablet, oral: 2.5 mg [BCF], 5 mg [BCF], 10 mg [BCF], 20 mg [BCF], 30 mg [BCF], 40 mg [BCF]   |
| Lisinopril and Hydrochlorothiazide | Tablet, oral:<br>10/12.5: Lisinopril 10 mg and hydrochlorothiazide 12.5 mg [BCF]<br>20/12.5: Lisinopril 20 mg and hydrochlorothiazide 12.5 mg [BCF]<br>20/25: Lisinopril 20 mg and hydrochlorothiazide 25 mg [BCF]   |
| Lithium                            | Capsule, oral, as carbonate: 150 mg [BCF], 300 mg [BCF]<br>Solution, oral, as citrate: 300 mg/5 mL (5 mL, 500 mL) [equivalent to amount of lithium in lithium carbonate] [BCF]<br>Tablet, extended release, oral, as carbonate: 450 mg<br>Lithobid: 300 mg   |
| Loperamide                         | Capsule, oral, as hydrochloride: 2 mg [BCF]  |
| Loratadine                         | Syrup, oral: 5 mg/5 mL (120 mL)<br>Tablet, oral: 10 mg   |
| Loratadine and Pseudoephedrine     | Tablet, extended release, oral: Loratadine 10 mg and pseudoephedrine sulfate 240 mg  |
| LORazepam                          | Tablet, oral: 0.5 mg, 1 mg   |
| Losartan                           | Tablet, oral, as potassium: 25 mg [BCF]<br>Cozaar: 50 mg [contains potassium 4.24 mg (0.108 mEq)] [BCF]<br>Cozaar: 100 mg [contains potassium 8.48 mg (0.216 mEq)] [BCF]   |
| Losartan and Hydrochlorothiazide   | Tablet, oral:<br>Hyzaar 50/12.5: Losartan potassium 50 mg and hydrochlorothiazide 12.5 mg [contains potassium 4.24 mg (0.108 mEq)] [BCF]<br>Hyzaar 100/12.5: Losartan potassium 100 mg and hydrochlorothiazide 12.5 mg [contains potassium 8.48 mg (0.216 mEq)] [BCF]<br>Hyzaar 100/25: Losartan potassium 100 mg and hydrochlorothiazide 25 mg [contains potassium 8.48 mg (0.216 mEq)] [BCF] |
| Loteprednol                        | Suspension, ophthalmic, as etabonate [drops]:<br>Alrex: 0.2% (10 mL) [contains benzalkonium chloride]  |
| Lurasidone                         | Tablet, Oral, as hydrochloride:<br>Latuda: 20 mg, 40 mg, 60 mg, 80 mg, 120 mg  |
| Magnesium Oxide                    | Tablet, oral:<br>MAGnesium-Oxide: 400 mg [equivalent to elemental magnesium 240 mg]  |
| Meclizine                          | Caplet, oral, as hydrochloride: 12.5 mg<br>Tablet, oral, as hydrochloride: 25 mg   |
| MedroxyPROGESTERone                | Injection, suspension, as acetate:<br>Depo-Provera Contraceptive: 150 mg/mL (1 mL) [contains polysorbate 80]<br>Tablet, oral, as acetate: 5 mg [BCF]<br>Provera: 2.5 mg [BCF], 10 mg [BCF] [scored]  |
| Megestrol                          | Tablet, oral, as acetate: 40 mg  |
| Meloxicam                          | Tablet, oral:<br>Mobic: 7.5 mg [BCF], 15 mg [BCF]  |
| Meperidine                         | Tablet, oral, as hydrochloride:<br>Demerol: 50 mg [scored]   |
| Mesalamine                         | Tablet, delayed release, enteric coated, oral:<br>Asacol: 400 mg   |
| MetFORMIN                          | Tablet, oral, as hydrochloride: 500 mg [BCF]<br>Glucophage: 850 mg [BCF]<br>Glucophage: 1000 mg [scored]<br>Tablet, extended release, oral, as hydrochloride: 500 mg [BCF], 750 mg [BCF]   |
| MethIMAzole                        | Tablet, oral:<br>Tapazole: 10 mg [scored]  |
| Methocarbamol                      | Tablet, oral: 500 mg [BCF], 750 mg   |
| Methotrexate                       | Injection, solution: 25 mg/mL (2 mL)<br>Tablet, oral: 2.5 mg [BCF]   |

Davis-Monthan AFB Outpatient Formulary

|                           |  |
|---------------------------|--|
| Methylphenidate           | <p>Capsule, extended release, oral, as hydrochloride:<br/>                     Metadate CD: 10 mg, 20 mg, 40 mg<br/>                     Ritalin LA: 10 mg [5 mg immediate release, 5 mg extended release] [BCF]<br/>                     Ritalin LA: 20 mg [10 mg immediate release, 10 mg extended release] [BCF]<br/>                     Ritalin LA: 30 mg [15 mg immediate release, 15 mg extended release] [BCF]<br/>                     Ritalin LA: 40 mg [20 mg immediate release, 20 mg extended release] [BCF]<br/>                     Suspension Reconstituted, Oral, as hydrochloride:<br/>                     Quillivant XR: 25 mg/5 mL<br/>                     Tablet, oral, as hydrochloride:<br/>                     Ritalin: 5 mg [BCF], 10 mg [scored] [BCF]<br/>                     Tablet, extended release, oral, as hydrochloride [bi-modal release]:<br/>                     Concerta: 18 mg [4 mg immediate release, 14 mg extended release] [BCF]<br/>                     Concerta: 27 mg [6 mg immediate release, 21 mg extended release] [BCF]<br/>                     Concerta: 36 mg [8 mg immediate release, 28 mg extended release] [BCF]<br/>                     Concerta: 54 mg [12 mg immediate release, 42 mg extended release] [BCF]<br/>                     Tablet, sustained release, oral, as hydrochloride: 20 mg</p> |
| MethylPREDNISolone        | Tablet, oral [dose-pack]: 4 mg (21s)   |
| Metoclopramide            | <p>Solution, oral: 5 mg/5 mL (473 mL) [BCF]<br/>                     Tablet, oral: 10 mg [BCF]</p>   |
| MetOLazone                | <p>Tablet, oral: 2.5 mg<br/>                     Zaroxolyn: 5 mg</p>   |
| Metoprolol                | <p>Tablet, oral, as tartrate: 25 mg [BCF], 50 mg [BCF], 100 mg [BCF]<br/>                     Tablet, extended release, oral, as succinate [expressed as mg equivalent to tartrate]: 25 mg [BCF], 50 mg [BCF], 100 mg [BCF]</p>  |
| MetroNIDAZOLE (Systemic)  | <p>Tablet, oral: 500 mg [BCF]<br/>                     Flagyl: 250 mg [BCF]</p>  |
| MetroNIDAZOLE (Topical)   | <p>Gel, topical:<br/>                     MetroGel: 1% (55 g)<br/>                     Gel, vaginal:<br/>                     MetroGel-Vaginal: 0.75% (70 g) [BCF]</p>   |
| Miconazole (Topical)      | <p>Cream, vaginal, as nitrate:<br/>                     Monistat 7: 2% (45 g) [contains benzoic acid; 100 mg/applicator]<br/>                     Powder, topical, as nitrate:<br/>                     Zeasorb-AF: 2% (70 g)</p>  |
| Minocycline (Systemic)    | Capsule, oral: 50 mg, 100 mg   |
| MiSOPROStol               | <p>Tablet, oral:<br/>                     Cytotec: 200 mcg [scored]</p>  |
| Modafinil                 | <p>Tablet, Oral:<br/>                     Provigil: 100 mg, 200 mg<br/>                     Generic: 100 mg, 200 mg</p>  |
| Montelukast               | <p>Tablet, oral:<br/>                     Singulair: 10 mg [contains lactose 89.3 mg/tablet] [BCF]<br/>                     Tablet, chewable, oral:<br/>                     Singulair: 4 mg [contains phenylalanine 0.67 mg/tablet; cherry flavor] [BCF]<br/>                     Singulair: 5 mg [contains phenylalanine 0.84 mg/tablet; cherry flavor] [BCF]</p>  |
| Morphine (Systemic)       | <p>Tablet, oral, as sulfate: 15 mg, 30 mg<br/>                     Tablet, controlled release, oral, as sulfate:<br/>                     MS Contin: 15 mg [BCF], 60 mg [BCF]<br/>                     Tablet, extended release, oral, as sulfate: 30 mg [BCF]</p>   |
| Moxifloxacin (Ophthalmic) | <p>Solution, ophthalmic [drops]:<br/>                     Vigamox: 0.5% (3 mL)</p>   |
| Moxifloxacin (Systemic)   | <p>Tablet, oral:<br/>                     Avelox: 400 mg</p>   |

Davis-Monthan AFB Outpatient Formulary

|  |   |
|--|---|
| Mupirocin  | Cream, topical, as calcium [strength expressed as base]:<br>Bactroban Cream: 2% (30 g) [contains benzyl alcohol]<br>Ointment, topical: 2% (22 g) [BCF]<br>Ointment, intranasal, as calcium [strength expressed as base]:<br>Bactroban Nasal: 2% (1 g)   |
| Mycophenolate  | Capsule, oral, as mofetil: 250 mg   |
| Naloxone   | Liquid, Nasal:<br>Narcan: 4 mg/0.1 mL (1 ea) [contains benzalkonium chloride, edetate disodium] [BCF]   |
| Naphazoline and Pheniramine                            | Solution, ophthalmic [drops]:<br>Naphcon A: Naphazoline hydrochloride 0.025% and pheniramine maleate 0.3% (5 mL)<br>[contains benzalkonium chloride]  |
| Naproxen   | Suspension, oral: 125 mg/5 mL (500 mL) [BCF]<br>Tablet, oral: 250 mg [BCF], 375 mg, 500 mg [BCF]<br>Tablet, oral, as sodium:<br>Anaprox: 275 mg [contains sodium 25 mg; equivalent to naproxen base 250 mg]<br>Anaprox DS: 550 mg [scored; contains sodium 50 mg; equivalent to naproxen base 500 mg] |
| Neomycin, Polymyxin B, and Dexamethasone               | Ointment, ophthalmic:<br>Maxitrol: Neomycin 3.5 mg, polymyxin B sulfate 10,000 units, and dexamethasone 0.1% per g (3.5 g)<br>Suspension, ophthalmic [drops]:<br>Maxitrol: Neomycin 3.5 mg, polymyxin B sulfate 10,000 units, and dexamethasone 0.1% per 1 mL (5 mL) [contains benzalkonium chloride] |
| Neomycin, Polymyxin B, and Gramicidin                  | Solution, ophthalmic [drops]:<br>Neosporin: Neomycin 1.75 mg, polymyxin B 10,000 units, and gramicidin 0.025 mg per 1 mL (10 mL) [BCF]  |
| Neomycin, Polymyxin B, and Hydrocortisone (Ophthalmic) | Suspension, ophthalmic [drops]:<br>Neomycin 3.5 mg, polymyxin B 10,000 units, and hydrocortisone 10 mg per 1 mL (7.5 mL)  |
| Neomycin, Polymyxin B, and Hydrocortisone (Otic)       | Solution, otic:<br>Neomycin 3.5 mg, polymyxin B 10,000 units, and hydrocortisone 10 mg per 1 mL (10 mL)<br>Suspension, otic:<br>Neomycin 3.5 mg, polymyxin B 10,000 units, and hydrocortisone 10 mg per 1 mL (10 mL)<br>[BCF]   |
| Nepafenac  | Suspension, ophthalmic [drops]:<br>Nevanac: 0.1% (3 mL) [contains benzalkonium chloride]  |
| Niacin   | Tablet, oral: 50 mg, 500 mg<br>Tablet, extended release, oral:<br>Niaspan: 500 mg [BCF], 750 mg [BCF], 1000 mg [BCF]  |
| Nicotine   | Patch, transdermal: 7 mg/24 hours (7s) [BCF] ; 14 mg/24 hours (14s) [BCF] ; 21 mg/24 hours (7s) [BCF]   |
| NIFEdipine   | Tablet, extended release, oral: 30 mg [BCF]<br>Adalat CC: 60 mg [BCF], 90 mg [BCF]  |
| Nitrofurantoin   | Capsule, oral [macrocrystal]: 50 mg [BCF]<br>Capsule, oral [macrocrystal/monohydrate]: 100 mg [BCF]<br>Suspension, oral: 25 mg/5 mL (230 mL)  |
| Nitroglycerin  | Patch, transdermal: 0.4 mg/hour (30s) [BCF]<br>Nitro-Dur: 0.2 mg/hour (30s) [BCF]<br>Solution, translingual [spray]:<br>Nitrolingual: 0.4 mg/spray (12 g) [contains ethanol 20%; 200 metered sprays] [BCF]<br>Tablet, sublingual:<br>Nitrostat: 0.4 mg [BCF]  |
| Norethindrone  | Tablet, oral:<br>Ortho Micronor: 0.35 mg [BCF]  |
| Nortriptyline  | Capsule, oral: 10 mg [BCF], 25 mg [BCF], 50 mg [BCF], 75 mg<br>Solution, oral: 10 mg/5 mL (473 mL) [BCF]  |

Davis-Monthan AFB Outpatient Formulary

|                             |   |
|-----------------------------|---|
| Nystatin and Triamcinolone  | Cream, topical: Nystatin 100,000 units and triamcinolone acetonide 0.1% (15 g)<br>Ointment, topical: Nystatin 100,000 units and triamcinolone acetonide 0.1% (15 g)   |
| Nystatin (Oral)             | Suspension, oral: 100,000 units/mL (60 mL, 473 mL) [BCF]  |
| Nystatin (Topical)          | Cream, topical: 100,000 units/g (15 g) [BCF]<br>Ointment, topical: 100,000 units/g (15 g) [BCF]   |
| Ofloxacin (Ophthalmic)      | Solution, ophthalmic [drops]: 0.3% (5 mL)   |
| Olopatadine (Nasal)         | Solution, intranasal [spray]:<br>Patanase: 0.6% (30.5 g) [contains benzalkonium chloride; equivalent to olopatadine hydrochloride 665 mcg/100 microliters; 240 metered sprays]  |
| Olopatadine (Ophthalmic)    | Solution, Ophthalmic [drops]:<br>Patanol: 0.1% (5 mL) [contains benzalkonium chloride] [BCF]<br>Pazeo: 0.7% (2.5 mL) [contains benzalkonium chloride]   |
| Omeprazole                  | Capsule, delayed release, oral: 20 mg [BCF]<br>PriLOSEC: 10 mg [BCF], 40 mg   |
| Ondansetron                 | Solution, oral: 4 mg/5 mL (50 mL)<br>Tablet, oral: 4 mg, 8 mg<br>Tablet, orally disintegrating, oral: 4 mg, 8 mg  |
| Oseltamivir                 | Capsule, oral, as phosphate:<br>Tamiflu: 75 mg<br>Powder for suspension, oral:<br>Tamiflu: 6 mg/mL (60 mL) [contains sodium benzoate; tutti frutti flavor]  |
| Oxybutynin                  | Tablet, Oral, as chloride:<br>Generic: 5 mg [BCF]<br>Tablet, extended release, oral, as chloride:<br>Ditropan XL: 5 mg [BCF], 10 mg [BCF], 15 mg [BCF]  |
| OxyCODONE                   | Tablet, oral, as hydrochloride: 5 mg  |
| Oxycodone and Acetaminophen | Tablet, oral:<br>Endocet: 5/325: Oxycodone hydrochloride 5 mg and acetaminophen 325 mg [scored] [BCF]   |
| Oxymetazoline (Nasal)       | Solution, intranasal, as hydrochloride [spray]: 0.05% (15 mL)   |
| Paliperidone                | Tablet Extended Release 24 Hour, Oral:<br>Generic: 1.5 mg, 3 mg, 6 mg, 9 mg   |
| Pancrelipase                | Capsule, delayed release, enteric coated microtablets [porcine derived]:<br>Pancrease: Lipase 21,000 units, protease 37,000 units, and amylase 61,000 units [ECF]   |
| Pantoprazole                | Tablet, delayed release, oral: 20 mg, 40 mg   |
| PARoxetine                  | Tablet, oral, as hydrochloride [strength expressed as base]: 10 mg, 20 mg, 30 mg, 40 mg   |
| Peginterferon Alfa-2a       | This medication is nonformulary. For formulary alternatives, refer to Therapeutic Category.<br>Kit, Subcutaneous [preservative free]:<br>Pegasys: 180 mcg/0.5 mL [contains benzyl alcohol] [ECF]<br>Solution, Subcutaneous [preservative free]:<br>Pegasys: 180 mcg/mL (1 mL) [contains benzyl alcohol] [ECF]<br>Pegasys: 180 mcg/0.5 mL (0.5 mL) [contains benzyl alcohol, polysorbate 80] [ECF] |
| PenicillAMINE               | Capsule, oral:<br>Cuprimine: 250 mg   |
| Penicillin V Potassium      | Powder for solution, oral: 250 mg/5 mL (200 mL) [BCF]<br>Tablet, oral: 250 mg [BCF], 500 mg [BCF]   |
| Pentoxifylline              | Tablet, controlled release, oral:<br>TRENtal: 400 mg  |
| Permethrin                  | Cream, topical: 5% (60 g) [BCF]   |
| Phenazopyridine             | Tablet, oral, as hydrochloride:<br>Pyridium: 100 mg [BCF]   |
| PHENobarbital               | Elixir, oral: 20 mg/5 mL (5 mL) [BCF]<br>Tablet, oral: 16.2 mg, 32.4 mg   |
| Phenylephrine (Nasal)       | Solution, intranasal, as hydrochloride [drops]:<br>Little Noses Decongestant: 0.125% (15 mL) [ethanol free; contains benzalkonium chloride]   |

Davis-Monthan AFB Outpatient Formulary

|  |   |
|--|---|
| Phenytoin                                | Capsule, extended release, oral, as sodium:<br>Dilantin: 100 mg [BCF]<br>Suspension, oral:<br>Dilantin-125: 125 mg/5 mL (240 mL) [contains ethanol 0.6%, sodium benzoate; orange-vanilla flavor] [BCF]<br>Tablet, chewable, oral:<br>Dilantin: 50 mg [scored] [BCF]   |
| Phytonadione                             | Tablet, oral:<br>Mephyton: 5 mg [scored]  |
| Pilocarpine (Ophthalmic)                 | Solution, ophthalmic, as hydrochloride [drops]: 2% (15 mL) [BCF]<br>Isopto Carpine: 1% (15 mL) [BCF]  |
| Pilocarpine (Systemic)                   | Tablet, oral, as hydrochloride:<br>Salagen: 5 mg  |
| Pimecrolimus                             | Cream, topical:<br>Elidel: 1% (30 g, 60 g, 100 g) [BCF] [contains benzyl alcohol]   |
| Pioglitazone                             | Tablet, oral:<br>Actos: 15 mg, 30 mg, 45 mg   |
| Piroxicam (Systemic)                     | Capsule, oral: 20 mg  |
| Podofilox                                | Gel, topical:<br>Condylox: 0.5% (3.5 g) [contains ethanol]  |
| Polyethylene Glycol 3350                 | Powder for solution, oral:<br>MiraLAX: 17 g/dose (510 g)  |
| Polyethylene Glycol-Electrolyte Solution | Powder for solution, oral: PEG 3350 240 g, sodium sulfate 22.72 g, sodium bicarbonate 6.72 g, sodium chloride 5.84 g, and potassium chloride 2.98 g (4000 mL)<br>TriLyte: PEG 3350 420 g, sodium bicarbonate 5.72 g, sodium chloride 11.2 g, and potassium chloride 1.48 g (4000 mL) [supplied with cherry, citrus berry, lemon lime, orange, and pineapple flavor packs] |
| Potassium Chloride                       | Capsule, extended release, microencapsulated, oral: 10 mEq [BCF]<br>Solution, oral: 20 mEq/15 mL (480 mL) [BCF]<br>Tablet, extended release, microencapsulated, oral: 10 mEq, 20 mEq [BCF]<br>Tablet, extended release, wax matrix, oral: 10 mEq [BCF]  |
| Potassium Citrate                        | Tablet, extended release, oral:<br>Urocit-K: 1080 mg [10 mEq]   |
| Pramipexole                              | Tablet, oral, as dihydrochloride monohydrate: 0.25 mg<br>Mirapex: 0.125 mg, 0.75 mg<br>Mirapex: 0.5 mg, 1 mg, 1.5 mg [scored]   |
| Pramoxine and Hydrocortisone             | Aerosol, foam, rectal:<br>Proctofoam HC: Pramoxine hydrochloride 1% and hydrocortisone acetate 1% (10 g)  |
| Pravastatin                              | Tablet, oral, as sodium: 10 mg [BCF]<br>Pravachol: 20 mg [BCF], 40 mg [BCF], 80 mg [BCF]  |
| Prazosin                                 | Capsule, oral: 1 mg, 2 mg, 5 mg   |
| PrednisoLONE (Ophthalmic)                | Suspension, ophthalmic, as acetate [drops]: 1% (5 mL) [BCF]   |
| PrednisoLONE (Systemic)                  | Solution, oral, as sodium phosphate [strength expressed as base]:<br>Orapred: 15 mg/5 mL (237 mL) [dye free; contains ethanol 2%, sodium benzoate; grape flavor] [BCF]  |
| PredniSONE                               | Tablet, oral: 1 mg [BCF], 5 mg [BCF], 10 mg [BCF], 20 mg [BCF]  |
| Primaquine                               | Tablet, oral, as phosphate: 26.3 mg [15 mg base]  |
| Primidone                                | Tablet, oral:<br>Mysoline: 50 mg, 250 mg [scored]   |
| Probenecid                               | Tablet, oral: 500 mg [BCF]  |
| Prochlorperazine                         | Tablet, oral, as maleate [strength expressed as base]: 5 mg, 10 mg  |
| Progesterone                             | Capsule, oral [micronized]:<br>Prometrium: 100 mg, 200 mg [contains peanut oil]   |

Davis-Monthan AFB Outpatient Formulary

|                                       |  |
|---------------------------------------|--|
| Promethazine                          | Suppository, rectal, as hydrochloride:<br>Promethegan: 25 mg (12s) [BCF]<br>Syrup, oral, as hydrochloride: 6.25 mg/5 mL (473 mL) [BCF]<br>Tablet, oral, as hydrochloride: 25 mg [BCF]  |
| Promethazine and Codeine              | Syrup, oral: Promethazine hydrochloride 6.25 mg and codeine phosphate 10 mg per 5 mL (118 mL)  |
| Promethazine and Dextromethorphan     | Syrup, oral: Promethazine hydrochloride 6.25 mg and dextromethorphan hydrobromide 15 mg per 5 mL (120 mL)  |
| Propafenone                           | Tablet, oral, as hydrochloride: 150 mg, 225 mg, 300 mg<br>Rythmol: 150 mg, 225 mg [scored]   |
| Propranolol                           | Capsule, extended release, oral, as hydrochloride: 60 mg<br>Capsule, sustained release, oral, as hydrochloride:<br>Inderal LA: 80 mg, 120 mg, 160 mg<br>Tablet, oral, as hydrochloride: 10 mg, 20 mg, 40 mg, 80 mg   |
| Propylthiouracil                      | Tablet, oral: 50 mg [BCF]  |
| Pseudoephedrine                       | Liquid, oral, as hydrochloride:<br>Children's Nasal Decongestant: 30 mg/5 mL (118 mL) [contains sodium benzoate; raspberry flavor]<br>Tablet, extended release, oral, as hydrochloride:<br>SudoGest 12 Hour: 120 mg  |
| Pyrazinamide                          | Tablet, oral: 500 mg [BCF]   |
| Pyridostigmine                        | Tablet, oral, as bromide:<br>Mestinon: 60 mg [scored]  |
| QUetiapine                            | Tablet, oral:<br>SEROquel: 25 mg [BCF], 50 mg [BCF], 100 mg [BCF], 200 mg [BCF], 300 mg [BCF]<br>Tablet, extended release, oral:<br>SEROquel XR: 50 mg [BCF], 150 mg [BCF], 200 mg [BCF], 300 mg [BCF], 400 mg [BCF]   |
| Raloxifene                            | Tablet, oral, as hydrochloride:<br>Evista: 60 mg [BCF]   |
| Ramipril                              | Capsule, oral:<br>Altace: 1.25 mg [BCF], 2.5 mg [BCF], 5 mg [BCF], 10 mg [BCF]   |
| RaNITidine (Withdrawn From US Market) | Syrup, oral:<br>Zantac: 15 mg/mL (480 mL) [contains ethanol 7.5%; peppermint flavor] [BCF]<br>Tablet, oral: 150 mg [BCF]   |
| Ribavirin (Systemic)                  | Capsule, Oral:<br>Generic: 200 mg [ECF]  |
| RifAMPin                              | Capsule, oral:<br>Rifadin: 150 mg, 300 mg [BCF]  |
| Rifapentine                           | Tablet, Oral:<br>Priftin: 150 mg   |
| RisperiDONE                           | Solution, oral: 1 mg/mL (30 mL) [BCF]<br>Tablet, oral:<br>RisperDAL: 0.25 mg [BCF], 0.5 mg [BCF], 1 mg [scored] [BCF], 2 mg [BCF], 3 mg [BCF], 4 mg [BCF]  |
| Rivaroxaban                           | Tablet, oral:<br>Xarelto: 10 mg, 15 mg, 20 mg  |
| Rizatriptan                           | Tablet, oral:<br>Maxalt: 5 mg [BCF], 10 mg [BCF]<br>Tablet, orally disintegrating:<br>Maxalt-MLT: 5 mg [contains phenylalanine 1.05 mg/tablet; peppermint flavor] [BCF]<br>Maxalt-MLT: 10 mg [contains phenylalanine 2.1 mg/tablet; peppermint flavor] [BCF] |
| ROPINIRole                            | Tablet, oral: 3 mg, 4 mg, 5 mg<br>Requip: 0.25 mg, 0.5 mg, 1 mg, 2 mg  |
| Rosuvastatin                          | Tablet, oral:<br>Crestor: 5 mg, 10 mg, 20 mg, 40 mg [Restricted: See Pharmacist]   |

Davis-Monthan AFB Outpatient Formulary

|  |  |
|--|--|
| Salicylic Acid   | Pad, topical:<br>Clear Away Wart Removing System: 40% (18s)  |
| Salmeterol   | Powder, for oral inhalation:<br>Serevent Diskus: 50 mcg (28s, 60s) [contains lactose] [BCF]  |
| Salsalate  | Tablet, oral: 500 mg [BCF], 750 mg [BCF]   |
| Selenium Sulfide   | Lotion, topical: 2.5% (120 mL) [BCF]   |
| Sertraline   | Tablet, oral: 25 mg [BCF], 50 mg [BCF], 100 mg [BCF]   |
| Sildenafil   | Tablet, oral:<br>Viagra: 25 mg [BCF], 50 mg [BCF], 100 mg [BCF]  |
| Silver Sulfadiazine                                      | Cream, topical:<br>Silvadene: 1% ( 400 g) [BCF]  |
| Simethicone  | Suspension, oral [drops]:<br>Mylicon Infants': 40 mg/0.6 mL (15 mL) [ethanol free; contains sodium benzoate]<br>Tablet, chewable, oral:<br>Mylanta Gas: 80 mg [peppermint flavor]  |
| Simvastatin  | Tablet, oral: 5 mg [BCF], 10 mg [BCF], 20 mg [BCF], 40 mg [BCF], 80 mg   |
| SITagliptin  | Tablet, oral:<br>Januvia: 25 mg [BCF], 50 mg [BCF], 100 mg [BCF]   |
| Sitagliptin and Metformin                                | Tablet, oral:<br>Janumet 50/500: Sitagliptin 50 mg and metformin hydrochloride 500 mg [BCF]<br>Janumet 50/1000: Sitagliptin 50 mg and metformin hydrochloride 1000 mg [BCF]<br>Tablet, extended release, oral:<br>Janumet XR: 50/500: Sitagliptin 50 mg and metformin hydrochloride 500 mg,<br>Janumet XR: 50/1000: Sitagliptin 50 mg and metformin hydrochloride 1000 mg,<br>Janumet XR:100/1000: Sitagliptin 100 mg and metformin hydrochloride 1000 mg  |
| Sodium Chloride  | Ointment, ophthalmic [preservative free]:<br>Muro 128: 5% (3.5 g)<br>Packet, Nasal [preservative free]:<br>Neilmed Saline Nasal Rinse: 1.57 g (100 ea)<br>Solution, for inhalation [preservative free]: 0.9% (5 mL)<br>Solution, intranasal [mist]:<br>Sea Soft Nasal Mist: 0.65% (45 mL)<br>Solution, ophthalmic [drops]:<br>Muro 128: 5% (15 mL)   |
| Sodium Phosphates  | Solution, rectal [enema]:<br>Fleet Enema: Monobasic sodium phosphate monohydrate 19 g and dibasic sodium phosphate heptahydrate 7 g per 118 mL delivered dose (133 mL) [contains sodium 4.4 g/118 mL]<br>Fleet Pedia-Lax Enema: Monobasic sodium phosphate monohydrate 9.5 g and dibasic sodium phosphate heptahydrate 3.5 g per 59 mL delivered dose (66 mL) [contains sodium 2.2 g/59 mL]<br>Tablet, oral:<br>OsmoPrep: Sodium phosphate monobasic monohydrate 1.102 g and sodium phosphate dibasic anhydrous 0.398 g [scored; gluten free; sodium phosphate 1.5 g per tablet] |
| Sodium Sulfate, Potassium Sulfate, and Magnesium Sulfate | Solution, oral:<br>Suprep Bowel Prep Kit: Sodium sulfate 17.5 g, potassium sulfate 3.13 g, and magnesium sulfate 1.6 g per 180 mL (180 mL) [contains sodium benzoate]  |
| Solifenacin  | Tablet, oral, as succinate:<br>VESicare: 5 mg, 10 mg   |
| Sotalol  | Tablet, oral, as hydrochloride:<br>Betapace: 80 mg [scored]  |
| Spironolactone   | Tablet, oral: 25 mg [BCF], 50 mg, 100 mg   |

Davis-Monthan AFB Outpatient Formulary

|                                     |  |
|-------------------------------------|--|
| Sucralfate                          | Suspension, Oral:<br>Carafate: 1 g/10 mL (420 mL) [contains methylparaben; cherry flavor]<br>Tablet, oral:<br>Carafate: 1 g [scored]   |
| Sulfacetamide and Prednisolone      | Ointment, ophthalmic:<br>Blephamide: Sulfacetamide sodium 10% and prednisolone acetate 0.2% (3.5 g)<br>Suspension, ophthalmic [drops]:<br>Blephamide: Sulfacetamide sodium 10% and prednisolone acetate 0.2% (5 mL) [contains benzalkonium chloride]   |
| Sulfacetamide (Ophthalmic)          | Solution, ophthalmic, as sodium [drops]: 10% (15 mL) [BCF]   |
| Sulfamethoxazole and Trimethoprim   | Suspension, oral: Sulfamethoxazole 200 mg and trimethoprim 40 mg per 5 mL (473 mL) [BCF]<br>Tablet, oral: Sulfamethoxazole 400 mg and trimethoprim 80 mg [BCF]<br>Tablet, double strength, oral: Sulamethoxazole 800 mg and trimethoprim 160 mg [BCF]  |
| SulfaSALazine                       | Tablet, oral: 500 mg [BCF]<br>Tablet, delayed release, enteric coated, oral:<br>Azulfidine EN-tabs: 500 mg [BCF]   |
| Sulfur and Sulfacetamide            | Cleanser, topical: Sulfur 5% and sulfacetamide sodium 10% (170 g)  |
| Sulindac                            | Tablet, oral: 150 mg, 200 mg   |
| SUMATriptan                         | Injection, solution, as succinate [strength expressed as base]:<br>Imitrex: 6 mg/0.5 mL (0.5 mL) [cartridge] [BCF]<br>Tablet, oral, as succinate [strength expressed as base]: 25 mg [BCF], 50 mg [BCF], 100 mg [BCF]  |
| Tacrolimus (Systemic)               | Capsule, Oral:<br>Prograf: 0.5 mg, 1 mg  |
| Tacrolimus (Topical)                | Ointment, External:<br>Generic: 0.03% (30 g, 60 g, 100 g) [BCF]; 0.1% (30 g, 60 g, 100 g) [BCF]  |
| Tamoxifen                           | Tablet, oral: 10 mg [BCF], 20 mg [BCF]   |
| Tamsulosin                          | Capsule, oral, as hydrochloride: 0.4 mg [BCF]  |
| Telmisartan                         | Tablet, Oral:<br>Generic: 20 mg, 40 mg, 80 mg  |
| Telmisartan and Hydrochlorothiazide | Tablet, Oral:<br>Micardis HCT: 40/12.5: Telmisartan 40 mg and hydrochlorothiazide 12.5 mg<br>Micardis HCT: 80/12.5: Telmisartan 80 mg and hydrochlorothiazide 12.5 mg<br>Micardis HCT: 80/25: Telmisartan 80 mg and hydrochlorothiazide 25 mg  |
| Temazepam                           | Capsule, oral: 15 mg [BCF], 30 mg [BCF]  |
| Terazosin                           | Capsule, oral: 1 mg [BCF], 2 mg [BCF], 5 mg [BCF], 10 mg [BCF]   |
| Terbinafine (Systemic)              | Tablet, oral: 250 mg   |
| Terconazole                         | Suppository, vaginal:<br>Terazol 3: 80 mg (3s) [contains coconut oil (may have trace amounts), palm kernel oil (may have trace amounts)]   |
| Testosterone                        | Gel, topical:<br>Fortesta: 10 mg/actuation (60 g) [contains ethanol; 0.5 g gel/actuation; 120 metered actuations] [BCF]<br>Injection, oil, as cypionate:<br>Depo-Testosterone: 200 mg/mL (1 mL) [contains benzyl alcohol, benzyl benzoate, cottonseed oil]<br>Patch 24 Hour, Transdermal:<br>Androderm: 2 mg/24 hr (60 ea); 4 mg/24 hr (30 ea) |
| Tetracycline (Systemic)             | Capsule, oral, as hydrochloride: 500 mg [BCF]  |
| Theophylline                        | Capsule, extended release, oral:<br>Theo-24: 300 mg [BCF], 400 mg [BCF] [24 hours]<br>Solution, oral: 80 mg/15 mL (15 mL, 473 mL) [BCF]<br>Tablet, extended release, oral: 200 mg [BCF], 300 mg [BCF], 400 mg [BCF]  |

Davis-Monthan AFB Outpatient Formulary

|                                 |   |
|---------------------------------|---|
| Timolol (Ophthalmic)            | Gel forming solution, ophthalmic, as maleate [strength expressed as base, drops]:<br>Timolol GFS: 0.25% (5 mL) [BCF]; 0.5% (5 mL) [BCF]<br>Solution, ophthalmic, as maleate [strength expressed as base, drops]: 0.25% (5 mL) [BCF];<br>0.5% (5 mL) [BCF]   |
| Tiotropium                      | Aerosol Solution, Inhalation:<br>Spiriva Respimat: 2.5 mcg/actuation (4 g)<br>Powder, for oral inhalation:<br>Spiriva HandiHaler: 18 mcg/capsule (30s) [contains lactose] [BCF]   |
| Tobramycin and Dexamethasone    | Ointment, ophthalmic:<br>TobraDex: Tobramycin 0.3% and dexamethasone 0.1% (3.5 g) [contains chlorobutanol]<br>Suspension, ophthalmic [drops]:<br>TobraDex: Tobramycin 0.3% and dexamethasone 0.1% (5 mL) [contains benzalkonium chloride]   |
| Tobramycin (Ophthalmic)         | Ointment, ophthalmic:<br>Tobrex: 0.3% (3.5 g) [contains chlorobutanol]<br>Solution, ophthalmic [drops]:<br>Tobrex: 0.3% (5 mL) [contains benzalkonium chloride]   |
| Tolterodine                     | Capsule, extended release, oral, as tartrate:<br>Detrol LA: 2 mg [BCF], 4 mg [BCF]  |
| Topiramate                      | Tablet, oral:<br>Topamax: 25 mg, 50 mg, 100 mg, 200 mg  |
| TraMADol                        | Tablet, oral, as hydrochloride: 50 mg [BCF]   |
| Tranexamic Acid                 | Tablet, Oral:<br>Generic: 650 mg  |
| TraZODone                       | Tablet, oral, as hydrochloride: 50 mg [BCF], 150 mg [BCF]   |
| Tretinoin (Topical)             | Cream, topical: 0.1% (20 g)<br>Retin-A: 0.025% (20 g [BCF]); 0.05% (20 g) [BCF]<br>Gel, topical:<br>Avita: 0.025% (20 g) [contains ethanol 83%]   |
| Triamcinolone (Topical)         | Aerosol, topical, as acetonide:<br>Kenalog: 0.2 mg/2-second spray (63 g) [contains dehydrated ethanol 10.3%]<br>Cream, topical, as acetonide: 0.1% (15 g, 80 g, 454 g) [BCF]; 0.5% (15 g)<br>Ointment, topical, as acetonide: 0.1% (15 g, 80 g, 454 g)<br>Paste, oral topical, as acetonide: 0.1% (5 g) |
| Trifluridine                    | Solution, ophthalmic [drops]:<br>Viroptic: 1% (7.5 mL)  |
| Trihexyphenidyl                 | Elixir, oral, as hydrochloride: 2 mg/5 mL (473 mL) [BCF]<br>Tablet, oral, as hydrochloride: 2 mg [BCF]  |
| Trimethobenzamide               | Capsule, oral, as hydrochloride:<br>Tigan: 300 mg   |
| Trimethoprim and Polymyxin B    | Solution, ophthalmic [drops]:<br>Polytrim: Trimethoprim 1 mg and polymyxin B sulfate 10,000 units per 1 mL (10 mL) [contains benzalkonium chloride] [BCF]   |
| Tripolidine and Pseudoephedrine | Tablet, oral:<br>Aprodine: Tripolidine hydrochloride 2.5 mg and pseudoephedrine hydrochloride 60 mg [scored]  |
| Tropicamide                     | Solution, ophthalmic [drops]:<br>Mydracyl: 1% (15 mL) [contains benzalkonium chloride]  |
| Ulipristal                      | Tablet, Oral, as acetate:<br>Ella: 30 mg [BCF]  |
| Urea                            | Lotion, topical:<br>Carmol 10: 10% (180 mL)   |
| Ursodiol                        | Capsule, oral:<br>Actigall: 300 mg  |

Davis-Monthan AFB Outpatient Formulary

|                                   |  |
|-----------------------------------|--|
| ValACYclovir                      | Caplet, oral:<br>Valtrex: 500 mg<br>Valtrex: 1 g [scored]  |
| Valproic Acid and Derivatives     | Capsule, sprinkle, oral:<br>Depakote Sprinkle: 125 mg [BCF] [strength expressed as valproic acid]<br>Syrup, oral, as valproate sodium:<br>Depakene: 250 mg/5 mL (473 mL) [strength expressed as valproic acid]<br>Tablet, delayed release, oral:<br>Depakote: 125 mg [BCF], 250 mg [BCF] [strength expressed as valproic acid]<br>Tablet, extended release, oral:<br>Depakote ER: 250 mg [BCF], 500 mg [BCF] [strength expressed as valproic acid]   |
| Valsartan                         | Tablet, Oral:<br>Diovan: 40 mg [scored] [BCF], 80 mg [BCF], 160 mg [BCF], 320 mg [BCF]<br>Generic: 40 mg [BCF], 80 mg [BCF], 160 mg [BCF], 320 mg [BCF]  |
| Valsartan and Hydrochlorothiazide | Tablet, Oral:<br>Diovan HCT: 80 mg/12.5 mg: Valsartan 80 mg and hydrochlorothiazide 12.5 mg [BCF]<br>Diovan HCT: 160 mg/12.5 mg: Valsartan 160 mg and hydrochlorothiazide 12.5 mg [BCF]<br>Diovan HCT: 160 mg/25 mg: Valsartan 160 mg and hydrochlorothiazide 25 mg [BCF]<br>Diovan HCT: 320 mg/12.5 mg: Valsartan 320 mg and hydrochlorothiazide 12.5 mg [BCF]<br>Diovan HCT: 320 mg/25 mg: Valsartan 320 mg and hydrochlorothiazide 25 mg [BCF]<br>Generic: 80 mg/12.5 mg: Valsartan 80 mg and hydrochlorothiazide 12.5 mg [BCF]<br>Generic: 160 mg/12.5 mg: Valsartan 160 mg and hydrochlorothiazide 12.5 mg [BCF]<br>Generic: 160 mg/25 mg: Valsartan 160 mg and hydrochlorothiazide 25 mg [BCF]<br>Generic: 320 mg/12.5 mg: Valsartan 320 mg and hydrochlorothiazide 12.5 mg [BCF]<br>Generic: 320 mg/25 mg: Valsartan 320 mg and hydrochlorothiazide 25 mg [BCF] |
| Varenicline                       | Combination package, oral [dose-pack]:<br>Chantix: Tablet: 0.5 mg (11s) [white tablets] and Tablet: 1 mg (42s) [light blue tablets]<br>Tablet, oral:<br>Chantix: 1 mg  |
| Venlafaxine                       | Capsule Extended Release 24 Hour, Oral:<br>Effexor XR: 37.5 mg [BCF], 75 mg [BCF], 150 mg [BCF]<br>Generic: 37.5 mg [BCF], 75 mg [BCF], 150 mg [BCF]<br>Tablet, Oral:<br>Generic: 25 mg [BCF], 37.5 mg [BCF], 50 mg [BCF], 75 mg [BCF], 100 mg [BCF]<br>Tablet Extended Release 24 Hour, Oral:<br>Generic: 225 mg [BCF]  |
| Verapamil                         | Tablet Extended Release, Oral, as hydrochloride:<br>Calan SR: 120 mg [BCF], 180 mg [scored] [BCF], 240 mg [scored] [BCF]<br>Generic: 120 mg [BCF], 180 mg [BCF], 240 mg [BCF]  |
| Vitamins (Multiple/Pediatric)     | Content varies depending on product used.<br>Solution, oral:<br>Poly-Vi-Sol: 35 mg/mL (50 mL)<br>Poly-Vi-Sol With Iron: 10 mg/mL (50 mL)   |
| Vitamins (Multiple/Prenatal)      | Content varies depending on product used.<br>Tablet, oral:<br>Prenatal Plus  |
| Warfarin                          | Tablet, oral, as sodium:<br>Coumadin: 1 mg [BCF], 2 mg [BCF], 2.5 mg [BCF], 3 mg [BCF], 4 mg [BCF], 5 mg [BCF], 6 mg [BCF], 7.5 mg [BCF], 10 mg [BCF]  |
| White Petrolatum                  | Gel: 28.35 g   |
| Witch Hazel                       | Pad, topical:<br>Tucks Anti-Itch: 50% (40s, 100s) [dye free; contains ethanol]   |
| Zaleplon                          | Capsule, oral:<br>Sonata: 5 mg, 10 mg [contains tartrazine]  |
| Zinc Oxide                        | Ointment, topical: 20% (30 g)  |

Davis-Monthan AFB Outpatient Formulary

|          |  |
|----------|--|
| Zolpidem | Tablet, oral, as tartrate: 5 mg [BCF], 10 mg [BCF]<br>Tablet, extended release, oral, as tartrate: 6.25 mg, 12.5 mg [Restricted: See Pharmacist] |
|----------|--|