

## Medication Guide

### Fentanyl transdermal system, CII

#### Rx Only

#### Fentanyl transdermal system is:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require daily around-the-clock, long-term treatment with an opioid, in people who are already regularly using opioid pain medicine, when other pain treatments such as non-opioid pain medicines or immediate-release opioid medicines do not treat your pain well enough or you cannot tolerate them.
- A long-acting (extended-release) opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.
- Not for use to treat pain that is not around-the-clock.

#### Important information about Fentanyl transdermal system:

- **Get emergency help right away if you use too much Fentanyl transdermal system (overdose).** When you first start taking Fentanyl transdermal system, when your dose is changed, or if you take too much (overdose), serious or life threatening breathing problems that can lead to death may occur.
- Never give anyone else your Fentanyl transdermal system. They could die from using it. Store Fentanyl transdermal system away from children and in a safe place to prevent stealing or abuse. Selling or giving away Fentanyl transdermal system is against the law.
- If the patch accidentally sticks to a family member while in close contact, take the patch off, wash the area with water, and get emergency help right away because an accidental exposure to Fentanyl transdermal system can lead to death or other serious medical problems.
- Proper disposal of Fentanyl transdermal system after use and for unused patches when no longer needed: Fold the sticky sides of the patch together and flush down the toilet. Do not put patches in a trash can.

#### Do not use Fentanyl transdermal system if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.

#### Before applying Fentanyl transdermal system, tell your healthcare provider if you have a history of:

- head injury, seizures
- liver, kidney, thyroid problems
- problems urinating
- pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.

#### Tell your healthcare provider if you:

- have a fever
- **are pregnant or planning to become pregnant.** Prolonged use of Fentanyl transdermal system during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **are breastfeeding.** Fentanyl transdermal system passes into breast milk and may harm your baby.
- are taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking Fentanyl transdermal system with certain other medicines can cause serious side effects that could lead to death.

#### When using Fentanyl transdermal system:

- Do not change your dose. Apply Fentanyl transdermal system exactly as prescribed by your healthcare provider.
- See the detailed Instructions for Use for information about how to apply and dispose of the Fentanyl transdermal system patch.
- Do not apply more than 1 patch at the same time unless your healthcare provider tells you to.
- You should wear the Fentanyl transdermal system patch continuously for 3 days, unless advised otherwise by your healthcare provider.
- **Call your healthcare provider if the dose you are using does not control your pain.**
- **Do not stop using Fentanyl transdermal system without talking to your healthcare provider.**

**While using Fentanyl transdermal system DO NOT:**

- Take hot baths or sunbathe, use hot tubs, saunas, heating pads, electric blankets, heated waterbeds, or tanning lamps, or engage in exercise that increases your body temperature. These can cause an overdose that can lead to death.
- Drive or operate heavy machinery, until you know how Fentanyl transdermal system affects you. Fentanyl transdermal system can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with Fentanyl transdermal system may cause you to overdose and die.

**The possible side effects of Fentanyl transdermal system are:**

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain, itching, redness, or rash where the patch is applied. Call your healthcare provider if you have any of these symptoms and they are severe.

**Get emergency medical help if you have:**

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue or throat, extreme drowsiness, light-headedness when changing positions, or you are feeling faint.

These are not all the possible side effects of Fentanyl transdermal system. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information go to [dailymed.nlm.nih.gov](http://dailymed.nlm.nih.gov).** For questions about fentanyl transdermal system, please call Upsher-Smith Laboratories, Inc. at 1-855-899-9180. If this is a medical emergency, please call 911.

Manufactured for:

UPSHER-SMITH LABORATORIES, INC., Maple Grove, MN 55369

This Medication Guide has been approved by the U.S. Food and Drug Administration.


Issued: April 2014

**INSTRUCTIONS FOR USE**

**FENTANYL TRANSDERMAL SYSTEM, CII**

**INSTRUCTIONS FOR APPLYING A FENTANYL TRANSDERMAL SYSTEM PATCH**

**Be sure that you read, understand, and follow these Instructions for Use before you use Fentanyl transdermal system. Talk to your healthcare provider or pharmacist if you have any questions.**



**Parts of the Fentanyl transdermal system patch:**

**Before applying Fentanyl transdermal system**

- Each Fentanyl transdermal system patch is sealed in its own protective pouch. Do not remove a Fentanyl transdermal system patch from the pouch until you are ready to use it.
- Do not use a Fentanyl transdermal system patch if the pouch seal is broken or the patch is cut, damaged or changed in any way.
- Fentanyl transdermal system patches are available in 5 different doses and patch sizes. Make sure you have the right dose patch or patches that have been prescribed for you.

## Applying a Fentanyl transdermal system patch

### 1. Skin areas where the Fentanyl transdermal system patch may be applied:

#### For adults:

- Put the patch on the chest, back, flank (sides of the waist), or upper arm in a place where there is no hair (See Figures 1–4).

#### For children (and adults with mental impairment):

- Put the patch on the upper back (See Figure 2).** This will lower the chances that the child will remove the patch and put it in their mouth.

#### For adults and children

- Do not** put a Fentanyl transdermal system patch on skin that is very oily, burned, broken out, cut, irritated, or damaged in any way.
- Avoid sensitive areas or those that move around a lot. If there is hair, **do not shave (shaving irritates the skin)**. Instead, clip hair as close to the skin as possible (See Figure 5).
- Talk to your healthcare provider if you have questions about skin application sites.**

### 2. Prepare to apply a Fentanyl transdermal system patch:

- Choose the time of day that is best for you to apply Fentanyl transdermal system. Change it at about the same time of day (3 days or 72 hours after you apply the patch) or as directed by your healthcare provider.
- Do not wear more than one Fentanyl transdermal system patch at a time unless your healthcare provider tells you to do so. Before applying a new Fentanyl transdermal system patch, remove the patch you have been wearing.
- Clean the skin area with clear water **only**. **Pat skin completely dry.** Do not use anything on the skin such as soaps, lotions, oils, or alcohol before the patch is applied.

3. **Open the pouch:** Fold and tear at slit, or cut at slit taking care not to cut the patch. Remove the Fentanyl transdermal system patch. Each Fentanyl transdermal system patch is sealed in its own protective pouch. Do not remove the Fentanyl transdermal system patch from the pouch until you are ready to use it (See Figure 6).

4. **Peel:** Peel off both parts of the protective liner from the patch. Each Fentanyl transdermal system patch has a clear plastic backing that can be peeled off in two pieces. This covers the sticky side of the patch. Carefully peel this backing off. Throw the clear plastic backing away. **Touch the sticky side of the Fentanyl transdermal system patch as little as possible** (See Figure 7).

5. **Press:** Press the patch onto the chosen skin site **with the palm of your hand and hold there for at least 30 seconds** (See Figure 8). Make sure it sticks well, especially at the edges.

- Fentanyl transdermal system may not stick to all patients. You need to check the patches often to make sure that they are sticking well to the skin.
- If the patch falls off right away after applying, throw it away and put a new one on at a different skin site. See the section below called "Disposing of a Fentanyl transdermal system patch".
- If you have a problem with the patch not sticking
  - Apply first aid tape only to the edges of the patch.
  - If you continue to have problems with the patch sticking, you may cover the patch with Bioclusive™. These are special see-through adhesive dressings. **Never cover a Fentanyl transdermal system patch with any other bandage or tape.** Remove the backing from the Bioclusive™ dressing and place it carefully over the Fentanyl transdermal system patch, smoothing it over the patch and your skin.
- If your patch falls off later, but before 3 days (72 hours) of use, dispose of properly. See the section below "Disposing of a Fentanyl transdermal system patch". Apply a new Fentanyl transdermal system patch on at a different skin site. Be sure to let your healthcare provider know that this has happened, and do not replace the new patch until 3 days (72 hours) after you put it on (or as directed by your healthcare provider).**



6. Wash your hands when you have finished applying a Fentanyl transdermal system patch.
7. Remove a Fentanyl transdermal system patch after wearing it for 3 days (72 hours). See the section below "Disposing of a Fentanyl transdermal system patch". Choose a **different** skin site to apply a new Fentanyl transdermal system patch. Repeat Steps 2 through 6 above when applying a new Fentanyl transdermal system patch.

**Do not apply the new patch to the same place as the last one.**

#### **Water and Fentanyl transdermal system**

- You can bathe, swim or shower while you are wearing a Fentanyl transdermal system patch. If the patch falls off before 3 days (72 hours) after application, dispose of properly. See the section below "Disposing of a Fentanyl transdermal system patch". Apply a new Fentanyl transdermal system patch on at a different skin site. Be sure to let your healthcare provider know that this has happened, and do not replace the new patch until 3 days (72 hours) after you put it on (or as directed by your healthcare provider).

#### **Disposing of a Fentanyl transdermal system patch**

- Fold the used Fentanyl transdermal system patch in half so that the sticky side sticks to itself (See Figure 9). **Flush the used Fentanyl transdermal system patch down the toilet right away (See Figure 10). A used Fentanyl transdermal system patch can be very dangerous for or lead to death in babies, children, pets, and adults who have not been prescribed Fentanyl transdermal system.**
- Throw away any Fentanyl transdermal system patches that are left over from your prescription as soon as they are no longer needed. Remove the leftover patches from their protective pouch and remove the protective liner. **Fold the patches in half with the sticky sides together, and flush the patches down the toilet.** Do not flush the pouch or the protective liner down the toilet. These items can be thrown away in a trashcan.

Figure 9



Figure 10



This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Rx Only

Manufactured for:

UPSHER-SMITH LABORATORIES, INC.

Maple Grove, MN 55369

© 2013 Upsher-Smith Laboratories, Inc.

Revised 0514

Bioclusive™ is a trademark of Ethicon, Inc.