

MEDICATION GUIDE

FLUOXETINE CAPSULES, USP

(floo ox' e teen)

10 mg and 20 mg

Read the Medication Guide that comes with fluoxetine capsules before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment. Talk with your healthcare provider if there is something you do not understand or you want to learn more about.

Fluoxetine capsules are in a class of drugs called selective serotonin reuptake inhibitors (SSRIs), which are often used for the treatment of depression and anxiety disorders. Although fluoxetine capsules are not a treatment for depression, it contains fluoxetine hydrochloride, the same active ingredient in some antidepressants.

What is the most important information I should know about fluoxetine capsules?

Fluoxetine capsules and other antidepressant medicines may cause serious side effects including:

1. Suicidal thoughts or actions:

- **Fluoxetine capsules and other antidepressant medicines may increase suicidal thoughts or actions** in some children, teenagers, or young adults within the **first few months of treatment or when the dose is changed.**
- Depression or other serious mental illnesses are the most important causes of suicidal thoughts or actions.
- Watch for these changes and call your healthcare provider right away if you notice:
 - o New or sudden changes in mood, behavior, actions, thoughts, or feelings, especially if severe.
 - o Pay particular attention to such changes when fluoxetine capsules are started or when the dose is changed.

Keep all follow-up visits with your healthcare provider and call between visits if you are worried about symptoms.

Call your healthcare provider right away if you have any of the following symptoms, or call 911 if an emergency, especially if they are new, worse, or worry you:

- attempts to commit suicide
- acting on dangerous impulses
- acting aggressive or violent
- thoughts about suicide or dying
- new or worse depression
- new or worse anxiety or panic attacks
- feeling agitated, restless, angry or irritable
- trouble sleeping
- an increase in activity or talking more than what is normal for you
- other unusual changes in behavior or mood

Call your healthcare provider right away if you have any of the following symptoms, or call 911 if an emergency. Fluoxetine capsules may be associated with these serious side effects:

2. Serotonin Syndrome. This condition can be life-threatening and may include:

- agitation, hallucinations, coma or other changes in mental status

- coordination problems or muscle twitching (overactive reflexes)
- racing heartbeat, high or low blood pressure
- sweating or fever
- nausea, vomiting, or diarrhea
- muscle rigidity
- dizziness
- flushing
- tremor
- seizures

3. Severe allergic reactions:

- trouble breathing
- swelling of the face, tongue, eyes or mouth
- rash, itchy welts (hives) or blisters, alone or with fever or joint pain

4. Manic episodes:

- greatly increased energy
- severe trouble sleeping
- racing thoughts
- reckless behavior
- unusually grand ideas
- excessive happiness or irritability
- talking more or faster than usual

5. Seizures or convulsions

6. Changes in appetite or weight. Children and adolescents should have height and weight monitored during treatment.

7. Abnormal bleeding: Fluoxetine capsules and other antidepressant medicines may increase your risk of bleeding or bruising, especially if you take the blood thinner warfarin (Coumadin^{®*}, Jantoven^{®*}), a non-steroidal anti-inflammatory drug (NSAID's, like ibuprofen or naproxen), or aspirin.

8. Visual Problems

- eye pain
- changes in vision
- swelling or redness in or around the eye

Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.

9. Low salt (sodium) levels in the blood. Elderly people may be at greater risk for this. Symptoms may include:

- headache
- weakness or feeling unsteady
- confusion, problems concentrating or thinking or memory problems

Do not stop fluoxetine capsules without first talking to your healthcare provider. Stopping fluoxetine capsules too quickly may cause serious symptoms including:

- anxiety, irritability, high or low mood, feeling restless or changes in sleep habits
- headache, sweating, nausea, dizziness
- electric shock-like sensations, shaking, confusion

What are fluoxetine capsules?

Fluoxetine capsules are a prescription medicine used to treat premenstrual dysphoric disorder (PMDD).

It is not known if fluoxetine capsules are safe and effective in children.

Talk with your healthcare provider if you do not think that your condition is getting better with fluoxetine capsules treatment.

Who should not take fluoxetine capsules?

Do not take fluoxetine capsules if you:

- take a Monoamine Oxidase Inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid.
 - Do not take an MAOI within 5 weeks of stopping fluoxetine capsules unless directed to do so by your healthcare provider.
 - Do not start fluoxetine capsules if you stopped taking an MAOI in the last 2 weeks unless directed to do so by your healthcare provider.

Women who take fluoxetine capsules close in time to an MAOI may have serious or even life-threatening side effects. Get medical help right away if you have any of these symptoms:

- high fever
- uncontrolled muscle spasms
- stiff muscles
- rapid changes in heart rate or blood pressure
- confusion
- loss of consciousness (pass out)
- **take Mellaril®* (thioridazine). Do not take Mellaril®* within 5 weeks of stopping fluoxetine capsules because this can cause serious heart rhythm problems or sudden death.**
- **take the antipsychotic medicine pimozide (Orap®*) because this can cause serious heart problems.**

What should I tell my healthcare provider before taking fluoxetine capsules?

Ask if you are not sure.

Before starting fluoxetine capsules, tell your healthcare provider if you:

- Are taking certain drugs or treatments such as:
 - Triptans used to treat migraine headache
 - Medicines used to treat mood, anxiety, psychotic or thought disorders, including tricyclics, lithium, bupropion, SSRIs, SNRIs, MAOIs or antipsychotics
 - Tramadol and fentanyl
 - Over-the-counter supplements such as tryptophan or St. John's Wort
 - Electroconvulsive therapy (ECT)
- have liver problems
- have kidney problems
- have heart problems
- have or had seizures or convulsions
- have bipolar disorder or mania
- have low sodium levels in your blood
- have a history of a stroke

- have high blood pressure
- have or had bleeding problems
- are pregnant or plan to become pregnant. It is not known if fluoxetine capsules will harm your unborn baby.
- are breastfeeding or plan to breastfeed. Some fluoxetine may pass into your breast milk. Talk to your healthcare provider about the best way to feed your baby while taking fluoxetine capsules.

Tell your healthcare provider about all the medicines that you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Fluoxetine capsules and some medicines may interact with each other, may not work as well, or may cause serious side effects.

Your healthcare provider or pharmacist can tell you if it is safe to take fluoxetine capsules with your other medicines. Do not start or stop any medicine while taking fluoxetine capsules without talking to your healthcare provider first.

If you take fluoxetine capsules, you should not take any other medicines that contain fluoxetine hydrochloride:

- Symbyax [®]*
- Prozac [®]*
- Prozac [®]* Weekly™

How should I take fluoxetine capsules?

- Take fluoxetine capsules exactly as prescribed. Your healthcare provider may need to change the dose of fluoxetine until it is the right dose for you.
- Talk to your healthcare provider if your symptoms do not get better.
- Fluoxetine capsules may be taken with or without food.
- If you take too much fluoxetine, call your healthcare provider or go to the nearest hospital emergency room right away.

What should I avoid while taking fluoxetine capsules?

Fluoxetine capsules can cause sleepiness or may affect your ability to make decisions, think clearly, or react quickly. You should not drive, operate heavy machinery, or do other dangerous activities until you know how fluoxetine capsules affects you. Do not drink alcohol while using fluoxetine capsules.

What are the possible side effects of fluoxetine capsules?

Fluoxetine capsules may cause serious side effects, including:

- See “What is the most important information I should know about fluoxetine capsules?”
- **Problems with blood sugar control.** Women who have diabetes and take fluoxetine capsules may have problems with low blood sugar while taking fluoxetine capsules. High blood sugar can happen when fluoxetine capsules are stopped. Your healthcare provider may need to change the dose of your diabetes medicines when you start or stop taking fluoxetine capsules.
- **Feeling anxious or trouble sleeping**

The most common side effects of fluoxetine capsules include:

-
- | | |
|-----------------------|-------------------------------|
| • headache | • trouble sleeping (insomnia) |
| • weakness (asthenia) | • dizziness |
| • pain | • feeling anxious or nervous |
| • accidental injury | • thinking differently |

- infection
 - flu symptoms
 - nausea
 - diarrhea
 - sexual problems
 - runny or stuffy nose (rhinitis)
 - sore throat (pharyngitis)
-

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of fluoxetine capsules. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store fluoxetine capsules?

- Store at 20° to 25°C (68° to 77°F).
- **Protect from light.**

Keep fluoxetine capsules and all medicines out of the reach of children.

General information about fluoxetine capsules

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use fluoxetine capsules for a condition for which it was not prescribed. Do not give fluoxetine capsules to other people, even if they have the same condition. It may harm them.

This Medication Guide summarizes the most important information about fluoxetine capsules. If you would like more information, talk with your healthcare provider. You may ask your healthcare provider or pharmacist for information about fluoxetine capsules that is written for healthcare professionals. For more information about fluoxetine capsules call Mylan Pharmaceuticals Inc. at 1-877-446-3679 (1-877-4-INFO-RX).

What are the ingredients in fluoxetine capsules?

Active ingredient: fluoxetine hydrochloride, USP

Inactive ingredients: colloidal silicon dioxide, magnesium stearate, pregelatinized starch (corn), and sodium lauryl sulfate. In addition, each of the empty gelatin capsules contain the following:

10 mg - FD&C Blue No. 1, FD&C Green No. 3, FD&C Yellow No. 6, gelatin, and titanium dioxide

20 mg - FD&C Red No. 40, gelatin, and titanium dioxide.

The imprinting ink contains the following: black iron oxide, D&C Yellow No. 10 Aluminum Lake, FD&C Blue No. 1 Aluminum Lake, FD&C Blue No. 2 Aluminum Lake, FD&C Red No. 40 Aluminum Lake, propylene glycol and shellac glaze.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

*The brands listed are trademarks of their respective owners.

Mylan Pharmaceuticals Inc.
Morgantown, WV 26505 U.S.A.

REVISED MARCH 2016
FLUXPD:R6