

Medication Guide

HYDROCODONE BITARTRATE (hye'' droe koe' done bye tar' trate)
AND ACETAMINOPHEN (a seet'' a min' oh fen)**TABLETS CII**

Hydrocodone bitartrate and acetaminophen tablets are:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage moderate to moderately severe pain severe enough to require an opioid analgesics , when other pain treatments such as non-opioid pain medicines do not treat your pain well enough or you cannot tolerate them.
- An opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.

Important information about hydrocodone bitartrate and acetaminophen tablets:

- **Get emergency help right away if you take too much hydrocodone bitartrate and acetaminophen tablets (overdose).** When you first start taking hydrocodone bitartrate and acetaminophen tablets, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur.
- Never give anyone else your hydrocodone bitartrate and acetaminophen tablets. They could die from taking it. Store hydrocodone bitartrate and acetaminophen tablets away from children and in a safe place to prevent stealing or abuse. Selling or giving away hydrocodone bitartrate and acetaminophen tablets is against the law.

Do not take hydrocodone bitartrate and acetaminophen tablets if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.
- previously exhibited hypersensitivity to hydrocodone or acetaminophen.
- known hypersensitivity to other opioids which may exhibit cross-sensitivity to hydrocodone.

Before taking hydrocodone bitartrate and acetaminophen tablets, tell your healthcare provider if you have a history of:

- head injury, seizures
- problems urinating
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.
- liver, kidney, thyroid problems
- pancreas or gallbladder problems

Tell your healthcare provider if you are:

- **pregnant or planning to become pregnant.** Prolonged use of hydrocodone bitartrate and acetaminophen tablets during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **Breast feeding.** Hydrocodone bitartrate and acetaminophen passes into breast milk and may harm your baby.

- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking hydrocodone bitartrate and acetaminophen tablets with certain other medicines can cause serious side effects that could lead to death.

When taking hydrocodone bitartrate and acetaminophen tablets:

- Do not change your dose. Take hydrocodone bitartrate and acetaminophen tablets exactly as prescribed by your healthcare provider.
- Take your prescribed dose every four to six hours at the same time every day. Do not take more than your prescribed dose. If you miss a dose, take your next dose at your usual time.
- Call your healthcare provider if the dose you are taking does not control your pain.
- If you have been taking hydrocodone bitartrate and acetaminophen tablets regularly, do not stop taking hydrocodone bitartrate and acetaminophen tablets without talking to your healthcare provider.
- After you stop taking hydrocodone bitartrate and acetaminophen tablets, the unused tablets should be disposed of by flushing down the toilet.

While taking hydrocodone bitartrate and acetaminophen tablets DO NOT:


- Drive or operate heavy machinery, until you know how hydrocodone bitartrate and acetaminophen tablets affects you. Hydrocodone bitartrate and acetaminophen tablets can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with hydrocodone bitartrate and acetaminophen tablets may cause you to overdose and die.

The possible side effects of hydrocodone bitartrate and acetaminophen tablets:

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of hydrocodone bitartrate and acetaminophen tablets. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.  For more information go to dailymed.nlm.nih.gov

Manufactured for: **QUALITEST PHARMACEUTICALS, Huntsville AL 35811**
call 1-800-444-4011

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Issued: 04/2016

R0 8183819